



Contents lists available at ScienceDirect

Blood Reviews

journal homepage: www.elsevier.com/locate/blre

REVIEW

Current concepts in the prevention of pathogen transmission via blood/plasma-derived products for bleeding disorders[☆]Giovanni Di Minno ^{a,*}, Carlo Federico Perno ^b, Andreas Tiede ^c, David Navarro ^d, Mariana Canaro ^e, Lutz Gürtler ^f, James W. Ironside ^g^a Dipartimento di Medicina Clinica e Chirurgia, Regional Reference Centre for Coagulation Disorders, Federico II University, Via S. Pansini 5, 80131 Naples, Italy^b Department of Experimental Medicine and Surgery, University of Rome Tor Vergata, Via Montpellier 1, 00133 Rome, Italy^c Department of Hematology, Hemostasis, Oncology and Stem Cell Transplantation, Hannover Medical School, Carl-Neuberg-Str. 1, D-30625 Hannover, Germany^d Department of Microbiology, Microbiology Service, Hospital Clínico Universitario, School of Medicine, University of Valencia, Av Blasco Ibáñez 17, 46010 Valencia, Spain^e Department of Hemostasis and Thrombosis, Son Espases University Hospital, Carretera de Valdemossa, 79, 07120 Palma de Mallorca, Spain^f Max von Pettenkofer Institute for Hygiene and Medical Microbiology, University of München, Pettenkofer Str 9a, 80336 Munich, Germany^g National Creutzfeldt-Jakob Disease Research and Surveillance Unit, School of Clinical Sciences, University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU, UK

ARTICLE INFO

Available online xxxx

Keywords:

Pathogen
Transfusion
Blood safety
Clotting
Haemophilia

ABSTRACT

The pathogen safety of blood/plasma-derived products has historically been a subject of significant concern to the medical community. Measures such as donor selection and blood screening have contributed to increase the safety of these products, but pathogen transmission does still occur. Reasons for this include lack of sensitivity/specifity of current screening methods, lack of reliable screening tests for some pathogens (e.g. prions) and the fact that many potentially harmful infectious agents are not routinely screened for. Methods for the purification/inactivation of blood/plasma-derived products have been developed in order to further reduce the residual risk, but low concentrations of pathogens do not necessarily imply a low level of risk for the patient and so the overall challenge of minimising risk remains. This review aims to discuss the variable level of pathogenic risk and describes the current screening methods used to prevent/detect the presence of pathogens in blood/plasma-derived products.

© 2015 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Contents

1. Introduction	0
2. Methodology	0
3. Infectious agents present in blood/plasma-derived products: lessons from the past and current concerns	0
3.1. Potential risk	0
4. Screening for pathogens	0
4.1. Blood donation screening: assays for key pathogens	0
4.1.1. HIV, HBV and HCV testing	0
4.1.2. HTLV-1 testing	0
4.1.3. Syphilis testing	0
4.2. Testing for emerging viral pathogens	0
4.2.1. West Nile virus	0
4.2.2. Chikungunya virus	0
4.2.3. Parvovirus B19	0
4.2.4. Hepatitis E virus	0
4.3. Prions	0
4.4. Interpretation of test results and practicalities of screening	0
4.4.1. Assay specificity and sensitivity	0
4.4.2. The need for multiple methods of screening	0
4.4.3. Infective dose	0

* Corresponding author. Tel.: +39 081 746 2060; fax: +39 081 5466152.

E-mail address: diminno@unina.it (G. Di Minno).

4.4.4. Patient characteristics	0
5. Other means of reducing and managing pathogen risk	0
5.1. Selection and screening of donors	0
5.2. Vaccination	0
5.3. Purification and inactivation techniques	0
5.4. Product choice	0
6. Barriers to a minimal risk approach	0
7. Current knowledge gaps and areas of unmet need	0
8. Conclusions	0
Practice points	0
Research agenda	0
Conflict of interest statement	0
Role of funding source	0
Acknowledgements	0
Appendix A. Supplementary data	0
References	0

1. Introduction

Acute bleeding episodes can arise either because of inherited bleeding disorders (e.g. haemophilia, von Willebrand disease), acquired deficiency of haemostatic components (e.g. due to infection, malignancy or autoimmune disease), trauma, surgery or as a result of infection with an organism that causes haemorrhagic disease (e.g. Ebola or Marburg virus) [1]. Various treatment options exist for preventing or treating acute bleeding episodes, including fresh-frozen plasma/cryoprecipitate, platelets and plasma-derived/recombinant clotting factor concentrates [2,3]. The use of blood-derived and recombinant haemostatic products has increased markedly over recent years, as exemplified by the global use of factor VIII products (Fig. 1) [4]. This increased use has been driven by improved availability of clotting factors, increased life expectancy of

people with bleeding disorders [5,6], increased use of prophylaxis for severe bleeding disorders [7,8] and decreased risk of transmission of infectious agents.

Historically, the risk of transmission of infectious agents via blood/plasma-derived products has been of great concern to the medical community. This risk has reduced dramatically since the implementation of stricter donation screening/donor selection procedures and improved purification procedures, but cannot be fully eradicated. Furthermore, the implementation of pathogen inactivation technology for blood/plasma-derived products has further reduced the risk of transmission of both known and emerging pathogens, although results can be variable according to the methods used [9,10]. However, it is important to note that patient risk is highly dependent on the circumstances under which blood products are collected, handled and used. In general, clinicians assess the level of risk associated with the use of blood/plasma-derived products by evaluating factors such as patient characteristics (e.g. age, immune status, geographical location, lifestyle) and the nature of the pathogen (e.g. physical characteristics, level of virulence, chronicity of infection, prevalence). The presence of a particular pathogen within blood/plasma-derived products may pose a significant threat to specific patient groups (e.g. the elderly or immunocompromised), while being of low risk to the general population (e.g. HEV).

While the clinical assessment of risk is based on a variety of factors, the virological assessment of risk is based solely on the presence or absence of pathogens. The presence of pathogens implies the possibility of infection, so only pathogen-free products can be described as entirely risk-free. Adopting the virological approach (i.e. discarding all products which may contain infectious agents) is effective in reducing the rate of pathogen transmission, but may result in the unnecessary wastage of blood/plasma-derived products.

Since recombinant clotting factors are not derived from blood or plasma, they present a minimal risk of pathogen transmission (particularly third generation factors, which have no contact with blood/plasma-derived components whatsoever) and can therefore be considered safer than using plasma-derived clotting factor concentrates [11]. However, there has been concern that the use of recombinant clotting factors may be associated with an increased rate of inhibitor formation in patients who regularly receive these products [12]; this is another factor that may influence the overall clinical assessment of patient risk.

Recombinant clotting factors are available for the treatment of haemophilia A (FVIII) and B (FIX) and factor VII/XIII deficiency, while plasma-derived concentrates are available for most other clotting factors (including von Willebrand factor, fibrinogen and the vitamin K dependent clotting factors). In some cases only fresh frozen plasma and cryoprecipitate are available for replacement therapy (e.g. factor V deficiency). Ideally, plasma and plasma-derived products would be completely replaced by recombinant products in order to minimise the risk of pathogen transmission; however, this is not always possible.




Fig. 1. Use of plasma-derived and recombinant factor VIII products in selected countries. Results are from Stonebraker et al. (1996–2006) [4] and WFH Annual Global Surveys (2007–2012) [196].

A recent article comprehensively reviewed the pathogen safety of plasma-derived and recombinant clotting factors [13].

This review examines the potential for transmission of infectious agents that might be present in blood/plasma-derived products used to treat haemostatic disorders. We focus on the identities of these agents and the screening procedures used for their detection, as well as the limitations of screening. Current unmet needs in the field of pathogen safety of blood/plasma-derived products are also discussed.

2. Methodology

The PubMed database was interrogated from 1 January 2000 to the present using the search strings 'bleeding disorders OR haemophilia' and 'pathogens AND blood safety'. The search terms 'virus', 'bacteria', 'haemorrhagic disorders', 'von Willebrand disease', 'FVII deficiency' and 'FXIII deficiency' were added to this search but did not yield any additional pertinent result. The bibliographies of reviews were also used to identify relevant references and individual searches were conducted for information on specific pathogens. Information and opinions were also provided by the authors.

3. Infectious agents present in blood/plasma-derived products: lessons from the past and current concerns

A large number of pathogenic agents (including viruses, protozoan parasites and prions) can be transmitted via blood/plasma-derived products and are capable of causing disease in humans (Table 1). The presence of viruses in plasma-derived products became a concern in the 1980s, when 60–70% of patients with severe haemophilia became infected with human immunodeficiency virus (HIV-1) [6]. This concern continued with the discovery that 80% of patients treated with plasma-derived products prior to 1992 had become infected with hepatitis C virus (HCV) [5]. Current donor selection and screening practices have improved our ability to detect or reduce the presence of pathogens in blood/plasma-derived products; for example, the residual risk of transfusion-transmitted infection (TTI) with HIV/HBV/HCV has fallen to near or less than 1 per million transfused units [14,15]. Despite this success, however, a residual risk still remains.

3.1. Potential risk

The pathogenic agents shown in Table 1 (and the Supplementary Appendix) do not form an exhaustive list. Many microorganisms that are normally non-pathogenic have the potential to cause disease when responding to changes in the biological environment, or when transfused to an immunosuppressed patient. In addition, there is still a risk that new and emerging pathogens may enter the blood supply (Table 2).

4. Screening for pathogens

The standard assays commonly used for blood screening are nucleic acid amplification technology (NAT) and immunoassays for detection of antibody and/or antigen. Immunoassays are frequently used for screening purposes as multiple samples can be processed together and they may yield semi-quantitative results. NAT assays allow earlier pathogen detection than with immunoassays, but they are also more costly and complex. Assay selection is generally determined by the level of accuracy/speed required, but factors such as the resources available (e.g. staff, infrastructure), assay complexity and cost considerations (e.g. consumables) must also be considered. Most assays for blood donation screening are mandatory (particularly in Europe and North America) and the World Health Organization (WHO) recommends that all whole blood (and blood which has been processed by apheresis) should undergo pathogen screening before it is used for clinical or manufacturing purposes. Screening for HIV-1, HIV-2, HBV, HCV and *Treponema pallidum*

subspecies *pallidum* (*T. pallidum*; the causative agent of syphilis) is strongly advised. The WHO and World Federation of Hemophilia (WFH) suggest that countries should carry out individual routine screening for further pathogens based on epidemiological information for their region e.g. HTLV-1 and *Trypanosoma cruzi* [16]. The WFH also acknowledges the positive impact of HIV, HBV and HCV screening on global blood safety and recommends that these screening tests be implemented whenever possible [2]. Details of the serological tests carried out on individual donor plasma and NAT testing of plasma mini-pools are shown in Tables 3 and 4 [17].

It is recommended that the minimum evaluated sensitivity/specificity level of any assay used for blood donation screening should be 99.5% or higher [16]. However, not all assays fulfil these criteria as sensitivity/specificity levels vary according to assay type and type of micro-organism being tested for (see Table 1 for details). Therefore, the general advice is to screen donated blood to as high a standard as possible [16].

4.1. Blood donation screening: assays for key pathogens

4.1.1. HIV, HBV and HCV testing

In the early stages of infection with HIV, HBV or HCV, viraemia occurs in the host's bloodstream at variable levels. Viral antigens may appear at the same time as DNA/RNA, but more often become detectable several weeks later. Specific antibodies are measurable 2 to 6 weeks after infection; the time between initial infection and the appearance of detectable parameters of infection (e.g. viral nucleic acid/antigens/antibodies) is known as the 'window period' [18–20].

4.1.1.1. HIV. When screening blood for the presence of HIV-1/HIV-2, the use of a combined antigen/antibody assay is advised (combined HIV p24 Ag and anti-HIV-1 + anti-HIV-2 antibodies) as it allows earlier detection of infection. The performance of NAT testing is mandatory in many countries and further reduces the window period (from around 20 to 11 days) [20–22].

4.1.1.2. HBV. The majority of diagnostic laboratories focus on the detection of HBsAg, which is the first detectable serological marker of infection. However, there is a risk that the HBsAg concentration may decline to undetectable levels during the course of infection, yielding a false negative result [23]. Screening for antibodies to HBc is the most conservative approach for identifying potentially exposed donors, as this identifies all individuals who have ever immunologically experienced any type of HBV infection (either current, chronic or resolved) and who may experience viral reactivation during their lifetime (particularly under conditions of immunosuppression). However, assays to measure HBc antibodies are relatively non-specific and do not always correlate with the presence of HBV virus in plasma [18]. Also, we cannot exclude the possibility that donors who test positive for anti-HBc do not have pulsed recurrences of virus replication, resulting in the presence of low levels of HBV-DNA in plasma. For these reasons, national transfusion services do not always routinely screen donations for anti-HBc. NAT assays can be carried out, but their use is restricted by potentially low levels of viral DNA [16,18]. The combined use of HBsAg screening tests and NAT assays has reduced the window period for detection of HBV infection from approximately 60 to 35.5 days [20,24]. Mutant HBV strains (escape variants) should also be considered, as they may occasionally escape serological detection (although most can be detected by NAT assays) [24,25]. These HBV-variants are rare, but therefore more likely to enter and contaminate the blood supply as they are more difficult to detect [25]. In summary, the screening of blood supplies for the presence of HBV is effective, but an optimal screening system has not yet been defined.

4.1.1.3. HCV. HCV (both recent and chronic infection) can be detected by screening blood for the presence of both HCV antigen and HCV antibody

Table 1

Blood-transmissible pathogens.

Transmissible agent	Geographic distribution (% of global population infected)	Testing procedure	Detection level in general clinical use	Minimum detection level possible (in development)	Testing sensitivity	Testing specificity
Prions						
vCJD	Rare [94]	Solid state binding matrix/immuno-assay [55,95]	N/A	N/A	≥71% [55,95]	≥98% [55,95]
Viruses						
Chikungunya virus	Widespread (prevalent in Africa, Asia and Latin-America) [96]	Antibody/NAT [96]	NAT: 40–350 copies/ml [97]	NAT: 10–100 copies/ml [98]	Antibody: varies according to disease stage and test type. Immuno-chromatographic test: 1.9–3.9% (acute infection) ELISA: 3.9% (acute), 84.1% (convalescent) NAT: RT-PCR – 88.5% (acute) [99]	Antibody: varies according to disease stage and test type. Immuno-chromatographic test: 92.5–95.0% (acute infection) ELISA: 92.5% (acute), 91.0% (convalescent) NAT: RT-PCR – 100% (acute) [99]
Cytomegalovirus (CMV)	Widespread [69]	Antigen/NAT	NAT: 446 copies/ml (whole blood) [100]	139 copies/ml (NAT; qPCR) [101]	Not available	Not available
Dengue virus	Widespread; 50% of global population at risk [102]	Antibody/NAT	NAT: 160–600 copies/ml (DENV-1, 3 and 4) 2000–6000 copies/ml (DENV-2) [103] based on 1 PFU = 1000–3000 RNA copies [104]	N/A: current main research focus is on developing serotype-specific tests	Antibody: 96–98% (IgM) [105] NAT: High for DENV-1, 3 and 4; lower for DENV-2 [103]	Antibody: 78–91% (IgM) [105] NAT: 100% [103]
Hepatitis B virus (HBV)	Widespread; 29% infected, 5% chronically infected [106]	Antibody/NAT	NAT: Reliable (≥99% detection) from 84 copies/ml [107] based on 1 IU/ml = 5.6 copies/ml [108]	NAT: 42–52 copies/ml (Ultrio); 4–10 copies/ml (Ampliscreen/UltraQual) [58]	Antibody: ≥90% (anti-HBc assay), 98% (anti-HBsAg assay) [109] NAT: 99.3% (Procleix Ultrio) [107]	Antibody: ≥78.9 (anti-HBc assay), up to 100% (anti-HBsAg assay) [109] NAT: 99.8% (Procleix Ultrio) [107]
Hepatitis C virus (HCV)	Widespread; 3% [110]	Antibody/NAT	NAT: 17 copies/ml (4-reaction kit format), 1664 copies/ml (primary pool format), 8806 copies/ml (master pool format), based on 1 IU/ml = 2.5 copies/ml [111,112]	NAT: 6.1 copies/ml (50% detection point, individual samples)	Antibody: 99% (anti-HCV-cAg) [113,114] NAT: 70.7–90.2% [111], 99.9% (Procleix Ultrio) [107]	Antibody: 99.2% (anti-HCV-cAg) [113,114] NAT: High [111], 98.1% (Procleix Ultrio) [107]
Hepatitis E virus (HEV)	Widespread [115]	Antibody/NAT	NAT: 5.9–114 copies/ml (MP-NAT), 47.3–226 copies/ml (ID-NAT) [116] based on 1 IU/ml = 1.25 copies/ml [117]	NAT: ≥5.8 copies/ml [118] based on 1 IU/ml = 1.25 copies/ml [117]	Antibody: 72–98% [119] NAT: High. Varies from 5.9 copies/ml (Real-Star HEV RT-PCR) to 114 copies/ml (Ceeram) for MP-NAT and from 47.3 copies/ml (Real-Star HEV RT-PCR) to 226 copies/ml (amplicube HEV RT-PCR) for ID-NAT [116]	Antibody: 78–96% [119] NAT: High [116]
Human Herpesvirus-8 (KSHV)	Widespread, with variable prevalence (≤5–50%); endemic in sub-Saharan Africa, parts of Asia and Oceania [120]	Antibody/NAT	NAT: 63 copies/ml (Viracor IBT) [121]	N/A	Antibody: 65–100% [122] NAT: High (63 copies/ml) [121]	Antibody: 96–99% [122] NAT: High – no cross reactivity detected against adenoviruses, BKV, CMV, EBV, HSV-1, HSV-2, HHV-6 variant A, HHV-6 variant B, HHV-7, JCV, parvovirus B19, SV-40 or VZV. [121]
Human immunodeficiency virus (HIV-1 and HIV-2)	Worldwide (0.5%); endemic in sub-Saharan Africa (5%) [123]	Antibody/NAT	NAT: 20–50 copies/ml [124]	N/A	Antibody: ≥99.9% [125] NAT: 99.9% (Procleix Ultrio) [107]	Antibody: 99.5–99.9% [125] NAT: 99.8% (Procleix Ultrio) [107]
Human T-lymphotropic virus 1 and 2 (HTLV-1/2)	HTLV-1: Endemic in Japan, Africa, South America and Caribbean	Antibody/NAT	Not specified: antibody detection kits routinely used [127,128]	NAT: HTLV-1–1.2 copies/reaction (i.e. 12–48 copies/ml, based on a reaction volume of 10–50 µl); HTLV-2–19	Antibody: 100% [130] NAT: 99.4% (HTLV-1 and 2) [131]	Antibody: ≥99.4% [130] NAT: 98.5% (HTLV-1 and 2) [131]

Table 1 (continued)

Transmissible agent	Geographic distribution (% of global population infected)	Testing procedure	Detection level in general clinical use	Minimum detection level possible (in development)	Testing sensitivity	Testing specificity
Parvovirus B-19	HTLV-2; Prevalent in Central/West Africa, the Americas and Europe [126] Widespread, variable prevalence (0.2–40% in European and Asian blood donors) [37]	Antibody/NAT	NAT: 668 copies/ml (based on 1 IU/ml = 3.34 copies/ml) [132,133]	NAT: 10 copies/reaction (i.e. 100–500 copies/ml, based on a reaction volume of 10–50 µl) [134]	Antibody: 89.1% [135] NAT: ≥93% [132]	Antibody: 99.4% [135] NAT: ≥97% [132]
Parvovirus 4	Widespread, variable prevalence (2–35%) [136–138]	NAT	NAT: 200–500 copies/ml [139,140]	NAT: 10 copies/reaction (i.e. 100–500 copies/ml, based on a reaction volume of 10–50 µl) [134]	Not available	Not available
Torque-tenovirus (TTV)	Present worldwide at variable prevalence (higher in developing countries) [141]	NAT	NAT: no general test – PCR used to confirm presence.	NAT: ≥100 copies/reaction (i.e. 1000–5000 copies/ml, based on a reaction volume of 10–50 µl) [142]	NAT: Up to 100% [142]	NAT: ≥96% [142]
West Nile virus (WNV)	Widespread (Africa, West Asia, Middle East, Europe and North and South America) [90]	Antibody/NAT	NAT: 100 copies/ml [143]	NAT: ≥9.8 copies/ml [144]	Antibody: 50% (IgM), 86% (IgG) [145] NAT: High [144]	Antibody: 95% (IgM), 69% (IgG) [145] NAT: High [144]
Bacteria						
<i>Treponema pallidum</i> subspecies <i>pallidum</i>	Widespread; (0.5% of global population infected) [146]	Antibody/NAT/dark ground microscopy DGM		NAT: 32,000 copies/ml [147]	NAT: 82% [148] Antibody: 78–88% (primary syphilis), 100% (secondary), 95–100% (latent), 71–96 (late) [149] DGM: 79–97% [147]	NAT: 95% [148] Antibody: 96–99% DGM: 77–100% [147,149]
<i>Yersinia enterocolitica</i> , <i>Salmonella enterica</i> , <i>Listeria monocytogenes</i> , <i>Coxiella burnetii</i> (only during outbreaks)	Rare, donor selection dependent	Not done		N/A	N/A	N/A
Protozoa						
<i>Babesia</i> spp	Widespread; predominantly in United States and (less frequently) Europe [150]	Antibody/NAT	Antibody: 10000–100000 parasites/ml (Giemsa staining) [151]	NAT: 5000–10000 parasites/ml [151]	Antibody: ≥97% [152] NAT: High [151]	Antibody: ≥97% [152] NAT: 100% [151]
<i>Leishmania</i> spp	Widespread; 1.3 million new cases per year [153]	Antibody/NAT	NAT: ≥10 parasites/ml (PCR; sensitivity varies according to method) [154]	NAT: ≥10 parasites/ml [154]	Antibody: 75–95% [155] NAT: 98.7% (kinetoplast DNA PCR), 91% (ITS1 PCR), 53.8% (SLME PCR) [156]	Antibody: 70–98% [155] NAT: 57.1% (kinetoplast DNA PCR), 100% (ITS1 PCR and SLME PCR) [156]
<i>Plasmodium</i> spp	Widespread (most common in sub-Saharan Africa). 30% at risk [157]	Antibody/NAT	Microscopy: 50000 parasites/ml [158]	NAT: <10 parasites/ml [158]	Antibody: ≥95% [159] NAT: 76.1–100% [158]	Antibody: ≥97.7% [159] NAT: 89.6–100% [158]
<i>Trypanosoma brucei</i> gambiense/rhodesiense	Regional in Africa, 10% at risk [160]	Direct microscopic visualisation/Antibody/NAT	NAT: <100 trypanosomes/ml [161]	NAT: 1–10 trypanosomes/ml [162]	Antibody: CATT – 87–98% [161] NAT: 88% [163]	Antibody: CATT – 95% [161] NAT: 99.2% [163]

(continued on next page)

Table 1 (continued)

Transmissible agent	Geographic distribution (% of global population infected)	Testing procedure	Detection level in general clinical use	Minimum detection level possible (in development)	Testing sensitivity	Testing specificity
<i>Trypanosoma cruzi</i>	0.1–0.2% (mostly Latin and South America) [164]	Antibody/NAT	NAT: ≥0.5 parasites/ml [165]	NAT: 0.05 parasites/ml [165]	Antibody: 75–100% (majority > 90%) [166] NAT: 83.3–94.4% [165]	Antibody: ≥97% [166] NAT: 85–95% [165]

In cases where no overall figures are available, specificity/sensitivity has been described as low, medium or high (as appropriate).

CATT, card agglutination test for Trypanosomiasis; ID-NAT, individual donation NAT; MP-NAT, mini-pool NAT; NAT, nucleic acid testing; PCR, polymerase chain reaction; PFU, plaque-forming unit.

(anti-HCV). Seroconversion occurs at approximately 6–8 weeks post-infection; however, steady improvements in screening technology (including the adoption of NAT assays) have reduced the window period to approximately 1–3 weeks [20,26]. As with HIV, NAT assays are more useful for detecting early infection, although the issue of low viral RNA concentration persists [27,28].

4.1.2. HTLV-1 testing

HTLV-1 and HTLV-2 are endemic in some regions, but very rare in others, and therefore screenings are conducted on a geographical or at-risk basis. The presence of virus is mostly inferred by the detection of virus-specific antibodies, using sensitive immunoassays [16].

4.1.3. Syphilis testing

In this instance, screening involves detection of non-specific, non-treponemal or specific treponemal antibodies. NAT assays are generally not used [29]. Specific antibody tests identify all individuals who have ever been exposed to this bacterium (and may continue to yield positive results for more than ten years following initial infection), while the non-specific tests (e.g. VDRL or cardiolipin tests) are primarily of use in identifying donors who may have an active infection. Since *T. pallidum* is heat-sensitive and cannot readily withstand extended storage at low temperatures, storage at 4 °C for more than three days is sufficient to render the pathogen non-infectious. However, blood components (e.g. platelets) that are stored at temperatures of around 20 °C do present a risk of *T. pallidum* persistence. Therefore screening for antibodies of this organism is recommended [16].

4.2. Testing for emerging viral pathogens

4.2.1. West Nile virus

Blood can be screened for further pathogens as appropriate, according to geographical location, seasonal activity of the vector and also patient risk factors. A current viral pathogen of interest is the mosquito-borne flavivirus West Nile virus (WNV), which was confirmed to have been transmitted via transfusion in 2002 [30]. An immediate screening policy was put in place in the USA in order to reduce the risk of further transmission. This policy included deferral of any individual displaying symptoms of infection, quarantine of plasma collected during periods of high mosquito activity (when WNV is most prevalent) and the rapid development/use of WNV-specific NAT and serological assays. These measures were highly effective and caused a significant reduction in the number of confirmed cases of WNV transfusion-related transmission. However, WNV outbreaks still occur within the Americas, indicating a potential need for seasonal blood screening for WNV [31]. WNV outbreaks have also occurred in Europe (including Italy and Greece), prompting the implementation of seasonal blood screening procedures in the affected regions of those countries [32].

4.2.2. Chikungunya virus

This is another mosquito-borne pathogen that could potentially pose a risk to transfusion safety, although to date reports of transfusion-related transmission of this virus are rare [33]. A mutated form of the Chikungunya virus has been responsible for several epidemics in the past decade, spreading to the Reunion Islands in the Indian Ocean (2005), Italy (2007) and the Caribbean area (2012/2013) [34–36]. The

Table 2
Emerging blood-transmissible pathogens.

Recently emerging pathogen	Pathogen structure/classification	Year of emergence	Mode of transmission	References
Bas Congo virus	Enveloped; rhabdovirus	2009	Direct contact (human–human and zoonotic), nosocomial	[167]
Huaiyangshan	Enveloped; bunyavirus	2010	Direct contact (human–human), vector-borne, blood-borne, airborne (potentially)	[168,169]
Bunyavirus				
Influenza H5N1	Enveloped; orthomyxovirus	2005	Airborne, direct contact (human–human and zoonotic)	[170–172]
Influenza H7N9	Enveloped; orthomyxovirus	2012	Direct contact (zoonotic and potentially human–human), airborne	[172,173]
Lujo virus	Enveloped; arenavirus	2008	Airborne, direct contact (human–human and zoonotic), nosocomial, blood-borne (potentially)	[174,175]
Marseillevirus ^a	Marseillevirus	2010	Blood-borne, faecal-oral (?)	[176,177]
MERS coronavirus	Enveloped; coronavirus	2013	Airborne, direct contact (human–human and zoonotic), nosocomial	[178,179]
SARS coronavirus	Enveloped; coronavirus	2003	Faecal/oral, airborne, direct contact (human–human and zoonotic), nosocomial	[180–182]
NJ polyomavirus	Non-enveloped polyomavirus	2014	Direct contact, saliva, blood?	[183]

^a Unconfirmed – may have been the result of laboratory contamination [184].

Table 3

Summarised table of serologic tests on individual donor plasma (WFH 2012) [17].

Plasma source	Serologic tests performed on individual donor plasma ^{a,b}
US paid apheresis (Talecris, Grifols, others)	Syphilis, HIV-1/2, HBsAg, HCVAb, ALT
United States, recovered, unpaid	Syphilis, HIV-1/2, HTLV-1&2, HBcAb, HBsAg, HCVAb, ALT
Baxter BioScience: United States, Austria, Germany, Sweden, Czech Republic, Switzerland, Norway	Syphilis, HIV-1/2, HBsAg, HCVAb
CSL Behring: Austria, Denmark, Germany, United States	Syphilis (US), HIV-1/2, HBsAg, HCVAb
Biostest: Austria, Belgium, Germany, United States, Switzerland	HIV-1/2, HBsAg, HCVAb
Intersero: Austria, Belgium, Germany	Syphilis, HIV-1/2, HBsAg, HCVAb, ALT
Germany unpaid	Syphilis, HIV-1/2, HBsAg, HCVAb, ALT
Octapharma: Sweden, Austria, Germany	Syphilis, HIV-1/2, HBcAb, HBsAg, HBCAb, HCVAb
American Community Blood Centers, unpaid (Octapharma)	Syphilis, HIV-1/2, HTLV-1&2, HBcAb = anti-HBc, HBsAg, HCVAb
Finnish Red Cross BS: Finland, unpaid	Syphilis, HIV-1/2, HBsAg, HCVAb. HTLV-1&2 1st donation.
Sanquin: The Netherlands	Syphilis, HIV-1/2, HTLV-1&2, HBsAg, HBcAb, HCVAb
LFB: France	Syphilis, HIV-1/2, HTLV-1&2, HBcAb, HBsAb, HBsAg, HCVAb
Grifols: Spain, Czech Republic, Slovakia	Syphilis, HIV-1/2, HBsAg, HCVAb
Kedrion: Italy	Syphilis, HIV-1/2, HBsAg, HCVAb, ALT
National Bioproducts Institute: South Africa	Syphilis, HIV-1/2, HBsAg, HCVAb
Australian Red Cross Blood Service	Syphilis, HIV-1/2, HTLV-1&2, HBsAg, HCVAb
New Zealand Blood Service	Syphilis, HIV-1/2, HBsAg, HCVAb. HTLV-1&2 (1st donation only)
Centre for Transfusion Medicine, Singapore	Syphilis, HIV-1/2, HBsAg, HCVAb
National Blood Center, Malaysia	Syphilis, HIV-1/2, HBsAg, HCVAb
Hong Kong Red Cross BTS	Syphilis, HIV-1/2, HTLV-1&2, HBsAb, HBsAg, HCVAb
Taiwan Blood Services Foundation	Syphilis, HIV-1/2, HTLV-1&2, HBsAb, HBsAg, HCVAb, ALT
Japan	Syphilis, HIV-1/2, HTLV-1, HBcAb, HBsAb, HBsAg, HCVAb, ALT, B-19 parvovirus
Korean Red Cross: South Korea	Syphilis, HIV-1/2, p-24 antigen, HBsAg, HCVAb, ALT
Shanghai RAAS Blood Products Co: China	Syphilis, HIV-1/2, HBsAg, HCVAb, ALT
Bio Products Laboratory, UK: US paid apheresis	Syphilis, HIV-1/2, HBsAg, HCVAb. HTLV-1&2, HBcAb (recovered only)

^a Serological targets for HIV detection include p24 antigen and anti-HIV-1/2 antibodies [185,186].

^b Nontreponemal tests are routinely used to screen blood for the presence of *Treponema pallidum* subspecies *pallidum* (*T. pallidum*, the causative agent of syphilis). These tests measure both proteins from the *Treponema* cell surface and IgG/IgM antiphospholipid antibodies which are produced by the host in response to cell damage in the early stages of infection [150].

virus may be detected in blood donors by NAT, which will help to reduce the level of transmission risk [35].

4.2.3. Parvovirus B19

There is also concern about the possibility of parvovirus B19 in the blood supply. B19 is prevalent worldwide, with seroprevalence in blood donors varying from between 0.2–1.3% in the USA, Europe and Africa and 25–40% in Asia [37]. The risk of parvovirus transmission is higher when units of blood are pooled (e.g. to create batches of clotting factor concentrates, albumin etc.) and so individuals with bleeding disorders are at a higher risk of infection. B19 DNA was detected in 26% of clotting factor concentrates in a recent German study [38], and another study found that populations receiving blood-derived products were 1.7 times more likely to display antibodies to B19 than populations who

had not received blood products [39]. B19 lacks a lipid envelope, which renders it highly resistant to some methods of pathogen inactivation [40]. Screening of blood donations for B19 DNA is not currently routine, but many manufacturers only process plasma that has been screened for the absence of B19 DNA in order to reduce the risk of transmission [41]. Given the prevalence of B19 in different populations, it is difficult to define the residual TTI risk of this virus. However, it is clear that a transmission risk does exist.

4.2.4. Hepatitis E virus

The potential presence of hepatitis E virus (HEV) in blood or blood-derived products is relevant. Currently there seems to be a discrepancy between the number of HEV-RNA positive blood donations in Europe (ranging from 1 in 1240 in Germany to 1 in 1761 donations in the Netherlands) [42], and the low number of confirmed cases of hepatitis E in blood transfusion recipients (one confirmed case in the UK [2006], one in France [2007] and two in Germany [2014]) [43–45]. This indicates that the subject of HEV infectivity and pathogenicity needs to be investigated further [46–49]. There is also a debate over the necessity of introducing screening blood for the presence of HEV; the virus is not currently screened for in the UK and other European countries, although one study of English donors found HEV to be widespread (1 in 2848 donations) within the donor population [50]. Since infection with this virus can be harmful to immunocompromised individuals, the potential need for introducing HEV screening should be considered [42].

The residual risk for TTI of HBV, HCV, HIV and HEV in selected countries is given in Table 5.

4.3. Prions

Despite recent advances in methods for the detection of prions, no single method has been developed as a screening test for blood, although several methods in animal models show great promise [51–53]. In humans, a protocol for the evaluation of a blood-based test for its suitability in the diagnosis of variant Creutzfeldt–Jakob disease (vCJD) has been established, but no test yet appears to satisfy the requirements of sensitivity and specificity [54]. The only report so far of a blood-based diagnostic test for vCJD claimed an assay sensitivity of 71.4% and a specificity of 100% in symptomatic patients and its potential applicability as a screening test to detect asymptomatic vCJD infection has recently been investigated [55]. There is currently no strategy for confirming a positive screening result, although the protein misfolding cyclical amplification technique has recently been shown to yield positive results in buffy coat/white blood cell samples from a small number of patients with vCJD [56].

4.4. Interpretation of test results and practicalities of screening

4.4.1. Assay specificity and sensitivity

Extensive use of blood donor selection and testing does not always guarantee a safe product. If the test is insufficiently sensitive, then false negatives may occur. Alternatively, tests which are not sufficiently specific (e.g. anti-HBc assay for HBV) may cause false positive results, leading to an unnecessary decrease in the number of clotting products available [16]. Torque Tenovirus (TTV) is an example where testing specificity has been an issue, as this virus exists in various divergent forms (23 distinct genotypes have been identified thus far) [57]. Since TTV is often detected in healthy individuals and is not associated with any particular disease, routine screening for this virus is not considered to be necessary; even a test with excellent sensitivity/specificity would not contribute to increase the level of safety of blood/plasma-derived products with regard to TTV.

Insufficient assay sensitivity remains a rare but potential problem when blood donations are screened during the window period of initial HBV, HCV or HIV infection. An increase in testing sensitivity threshold is

Table 4

Plasma inventory hold and NAT testing of mini-pools (WFH 2012) [17].

Company or Fractionator	Mini-Pool NAT Tests ^a	Manufacturing Pool NAT Tests ^b	NAT on Final Product ^c	Inventory Hold ^d	Mini-Pool Size ^e
CSL Behring: United States, Germany	HAV, HBV, HCV, HIV-1, B-19 parvovirus	HAV, HBV, HCV, HIV, B-19 parvovirus	No	60+ days	512 or fewer
Baxter BioScience: United States, Austria, Italy	HAV, HBV, HCV, HIV-1, B-19 parvovirus	HAV, HBV, HCV, HIV-1/2, B-19 parvovirus	No	60+ days	512 or fewer
Talecris: United States	HBV, HCV, HIV-1, B-19 parvovirus	HBV, HCV, HIV-1, B-19 parvovirus	No	60+ days	96 or 480
Grifols: United States, Spain, Czech Republic, Slovakia	HAV, HBV, HCV, HIV, B-19 parvovirus	HBV, HCV, HIV, B-19 parvovirus		60+ days	512 or fewer
Bio Products Laboratory, UK	HAV, HBV, HCV, HIV-1/2, B-19 parvovirus	HCV		60 days	512 or fewer
Biotest: Germany	HAV, HBV, HCV, HIV-1/2, B-19 parvovirus	HBV, HCV, HIV	No	60 days	960
Intersero, Germany	HAV, HBV, HCV, HIV 1, B-19 parvovirus	HBV, HCV, HIV		60+ days	960
German Red Cross BSO NSTOB	HAV, HBV, HCV, HIV 1, B-19 parvovirus	HCV		2 months	48
Octapharma: Sweden, Austria, Germany, USA	HIV, B-19 parvovirus, HAV, HCV, HIV-1	HCV	No	2 months	16-512
Finnish Red Cross BS: Finland	HIV, HCV, HIV (individual)	FRC BS does not make plasma pools			1 or 96
Sanquin: The Netherlands	HAV, B-19 parvovirus (mini-pool) HCV (6), HIV (6), HBV (6), B-19 parvovirus (480), HAV (480)	HBV, HCV, HIV, B-19 parvovirus	No		480 or 6
LFB: France	1: B-19 parvovirus 2: HAV, HCV	HAV, HBV, HCV, HIV-1, B-19 parvovirus		80+ days	1: 300 2: 1000
Kedrion: Italy	HBV, HCV, HIV 1, B-19 parvovirus (HAV if required)	HCV		60+ days	480 or fewer
National Bioproducts Institute: South Africa	HAV, HCV, HIV, B-19 parvovirus	HAV, HCV, HIV			1 and 216
CSL Biotherapies: Australia	HCV, HIV	HCV, HIV, B-19 parvovirus			480
Australian Red Cross Blood Service	HCV, HIV, B-19 parvovirus (optional)	HCV, HIV, B-19 parvovirus			480/512
Fractionated at CSL Biotherapies					
New Zealand Blood Service Fractionated at CSL Biotherapies	HCV, HIV, B-19 parvovirus (optional)	HCV, HIV, B-19 parvovirus			480/512
Hong Kong Red Cross BTS Fractionated at CSL Biotherapies		HCV, HIV			
Blood Services Group, Singapore Fractionated at CSL Biotherapies	HCV, HIV	HCV, HIV			480/512
National Blood Centre of Malaysia	HCV, HIV	HCV, HIV			480/512
Fractionated at CSL Biotherapies					
Taiwan Blood Services Foundation	HAV, HBV, HCV, HIV, B-19 parvovirus	HCV, HIV (optional HBV, HAV, B-19 parvovirus)			480/512
Fractionated at CSL Biotherapies					
GreenCross: South Korea	HAV, HCV	HAV, HBV, HCV, HIV	HAV, HBV, HCV, HIV	45 days	<450
Japanese Red Cross: Japan	HBV, HCV, HIV-1	HBV, HCV, HIV-1	HAV, HBV, HCV, HIV-1, B-19 parvovirus	6 months	20
Kaketsukan: Japan	1: HBV, HCV, HIV-1 2: HAV, B-19 parvovirus	HAV, HBV, HCV, HIV-1, B-19 parvovirus	HAV, HBV, HCV, HIV-1, B-19 parvovirus	6 months	1: 50 2: 500
Benesis: Japan	HBV, HCV, HIV-1	HBV, HCV, HIV-1	HAV, HBV, HCV, HIV-1, B-19 parvovirus	6 months	50
Shanghai RAAS Blood Products: China	HBV, HCV, HIV-1	HBV, HCV, HIV-1	HAV, HBV, HCV, HIV-1	60+ days	48

HAV, hepatitis A virus; HBV, hepatitis B virus; HCV, hepatitis C virus; HIV-1, human immunodeficiency virus-1; NAT, nucleic acid testing.

^a The viruses which are screened for by the company or fractionator using NAT at mini-pool stage. Numbers in brackets indicate the mini-pool sizes for each NAT type (if applicable).^b The viruses which are screened for by the company or fractionator using NAT at manufacturing pool stage.^c The viruses which are screened for by the company or fractionator using NAT within the final product (if any).^d Length of time that plasma is retained between donation and processing stages for donor information gathering.^e A mini-pool is a pool of donor samples, formed by directly pooling samples from individual donors or by pooling of samples from subpools. The numbers in the table indicate the number of samples present in the pool [187].

needed to prevent HBV transmission via blood/plasma-derived products and by blood transfusion, as extremely low concentrations of HBV (e.g. 1.6 copies/ml) is capable of viral transmission (see above section on anti-HBc positive patients) [58]. In the case of HIV and HCV, even NAT testing may not always be sufficient to ensure sufficiently high levels of safety as virus transmission has previously occurred after transfusion of blood with undetectable levels of viraemia [59].

Recent reports have highlighted concerns about the inability of NAT assays to detect different variants of HIV. There have been at least four cases in which the presence of HIV-1 RNA was undetected by NAT assay screening, potentially putting transfusion recipients at risk [21]. Two of these false-negative results occurred due to genetic mutation in the viral RNA regions targeted by NAT assay primers (although in the majority of cases serology yielded a positive result) [60,61]. False-negative results can be avoided by designing NAT assays that target a minimum of two amplification regions; such testing will be mandatory in Germany from 2015 [21,62]. It is hoped that as the sensitivity and

specificity of NAT tests continue to improve, cases of undetected infection may become less of an issue in the future.

4.4.2. The need for multiple methods of screening

A recent report suggests that it is not necessary to carry out serological screening for multiple HBV markers, and that NAT based screening is preferred [58]. However, there is a risk that the practice of relying upon a single method of screening may lead to a higher incidence of false-negative results. In one case of transfusion-associated parvovirus B19 transmission, donated blood was screened for the presence of B19 antigen and deemed to be safe since no antigen was detected. Since the recipient subsequently developed B19 infection, the donation was re-analysed and found to contain B19 DNA [63]. Reports such as this support the argument for multiple parameter testing.

The localisation of pathogens within blood may also influence ease of detection. For example, WNV is present at 10-fold higher levels in whole blood than in plasma in viraemic seropositive donors. The

Table 5

Approximate residual risk of HBV, HCV, HIV and HEV TTI in selected countries.

Virus	Country	Residual risk, or prevalence in blood donations (dependent on testing) per 100,000	Reference
HBV	Congo-Kinshasa	390	[188]
	Brazil	289	[189]
	USA	5.9–7.5	[58]
	England/North Wales	<3.7	[190]
	Australia	2	[191]
	Germany	0.4	[192]
	Portugal	0.2	[193]
HCV	Brazil	191	[189]
	Germany	0.9	[192]
	Portugal	0.03	[193]
HIV	Congo-Kinshasa	22	[188]
	Germany	0.9–2.4	[21,192]
HEV	Germany	81	[42]
	Netherlands	57	[42]
	England	35	[50]
	Scotland	6.9	[194]
	Portugal	0.6	[193]
	USA	<2	[195]

HBV, hepatitis B virus; HCV, hepatitis C virus; HIV-1, human immunodeficiency virus-1; TTI, transfusion-transmitted infection.

situation is reversed in viraemic seronegative donors, who display higher WNV levels (4-fold) in plasma than in whole blood [64]. Also, cell-associated viruses such as CMV and HTLV-1 are less frequently transmitted with the use of leukoreduced products [65,66], indicating that these viruses are less likely to be found in blood/plasma-derived products. The minimum infective dose (MID: the lowest number of pathogenic particles required to successfully infect a host) of a particular pathogen is more likely to be reached in whole blood or blood-derived components, making the use of plasma-derived or recombinant clotting factors the safest option.

4.4.3. Infective dose

To date, studies attempting to measure human MID values have generally determined the viral concentration needed to infect a particular percentage of the exposed population (e.g. 50%). This value (the human infective dose for 50% of the population) is referred to as HID₅₀ and is often described as the human MID [67]. The HID₅₀ value varies greatly between pathogens (even if they are physically similar) [68] and also varies depending upon the immune status of the recipient, as immunocompromised individuals, neonates and the elderly are at greater risk of infection than healthy individuals [67]. When this finding and the prevalence of immunocompromised patients receiving blood products are both taken into account, it implies a need for screening tests to have the highest sensitivity possible.

4.4.4. Patient characteristics

The impact of transmission on morbidity and mortality is dependent on patient characteristics. For example, although the vast majority of cases of CMV infection are not clinically important, influencing factors such as genetic predisposition, malnutrition and pre-existing infection can lead to the development of severe disease [69]. National screening programmes do not currently screen for CMV as standard, but NAT assays exist and may be carried out if required, e.g. if blood is specifically intended for vulnerable recipients, such as pregnant women or transplant patients [70]. It is the opinion of the authors that both serological assays and NAT tests should be used in order to reach the highest level of safety possible, particularly when in the case of immunosuppressed patients.

5. Other means of reducing and managing pathogen risk

As well as blood donation testing, a range of other measures are used to increase the pathogen safety of blood- and plasma-derived products.

These include donor selection and screening, recipient vaccination and the use of blood product purification/inactivation methods. The choice of inactivation method also impacts upon the level of risk.

5.1. Selection and screening of donors

Questionnaires are often used to attempt to assess donors' health status and their potential exposure to various risks. Donors can be accepted or rejected on the basis of these answered questionnaires, or alternatively their blood may be put through additional screening tests as appropriate [16,31].

5.2. Vaccination

Patients with bleeding disorders should be vaccinated against HAV and HBV. European studies have reported that universal HBV vaccination of blood donors could be cost-effective as this measure would reduce the risk of HBV transmission in general and might even remove the necessity for general HBV NAT testing; however, this would not reduce the risk posed by HBV escape variants (as described earlier) [71,72].

5.3. Purification and inactivation techniques

Blood/plasma-derived products typically undergo various procedures designed to reduce the pathogen level, although these procedures are not effective against all pathogens. Plasma donations undergo quarantine (approximately 4–6 months) prior to fractionation and when the donor is again screened negative FFP can then be subjected to chromatographic fractionation, solvent-detergent treatment, nanofiltration and/or heat inactivation [73,74]. Prolongation of product storage time can be effective in reducing the infectivity of temperature-sensitive pathogens (such as *T. pallidum*). Production of recombinant products also follows strict protocols to remove and inactivate any viruses that might be present, even though the risk of viral presence is minimal.

Although current purification/inactivation techniques (such as solvent-detergent treatment, nanofiltration and heat activation) do reduce the risk of pathogen transmission, they are not always sufficient to render blood/plasma-derived products safe [75]. Small non-enveloped viruses (e.g. HAV, B19 and picornavirus) are often highly resistant to inactivation procedures and may still be infectious in some plasma-derived concentrates [75,76].

The presence of prions is also a concern. Attempts to remove prions from plasma-derived products have involved several techniques, including ion-exchange chromatography and nanofiltration [77,78]. Several problems exist with these approaches, in particular the use of exogenous "spikes" derived from prion-infected brain homogenates to measure prion clearance, which may result in an overestimation of the amount of prion removal, and the methods used for the estimation of the reduction in prion load, which ideally should involve bioassay to measure infectivity [79]. Further developments in this field are required to address these issues.

A recently proposed approach for the inactivation of infectious agents in blood is whole-blood treatment with ultraviolet (UV) light in combination with a photosensitiser such as riboflavin or amotosalen [80,81]. The main disadvantage of this approach is that UV treatment has been linked to the formation of neoantigens, which may be generated via modification of the surface antigens of platelets. The presence of neoantigens may provoke a recipient immune response during transfusion of UV-treated platelets, causing them to be rapidly cleared from the circulation [81]. While this approach to pathogen inactivation is currently used for platelets and is effective, it needs further refinement for the inactivation of whole blood [81,82].

5.4. Product choice

The WFH strongly recommends the use of plasma-derived or recombinant products in preference to cryoprecipitate or fresh frozen plasma, as the infectious load of any infectious human pathogen is lower in plasma-derived products than in cryoprecipitate, and even lower in recombinant products [2]. In some European countries, recombinant products have almost completely replaced plasma-derived products [12]. The use of recombinant products, which have been manufactured and formulated with minimal addition of human/animal-derived materials, greatly reduces the risk of recipient exposure to plasma-derived infectious agents and after they have undergone virus removal/inactivation processes, recombinant products can be considered to be as safe as currently possible [83]. Despite these benefits, the use of recombinant products may be limited by higher costs and perceived problems of inhibitor formation [83]. For indications where no recombinant factor concentrates are available, the use of inactivated plasma-derived concentrates is safer than fresh frozen plasma and will reduce the risk of other adverse effects such as hypervolemia, transfusion-related lung injury (TRALI) and hypersensitivity [84].

6. Barriers to a minimal risk approach

A minimal risk approach would ensure that patients receive effective treatment with the lowest possible risk, but this is difficult to achieve in practical terms. Regulatory needs in different European countries are usually based on recommendations from the medical community, so in order to achieve minimal risk, it would be ideal for these regulations to be standardised and mandatory in all countries. Directives issued by the European Union Commission describe the regulatory requirements for the safety of whole blood and plasma, stating that "all precautionary measures during their collection, processing, distribution and use need to be taken making appropriate use of scientific progress in the detection and inactivation and elimination of transfusion transmissible pathogenic agents" [85]. However, the relative safety of different screening tests, products and processing methods is not discussed and so individual countries may adopt different approaches towards minimising risk. Although recombinant products are associated with the highest level of pathogen safety, higher costs for development and production may make them too expensive for some healthcare systems [86]. Inhibitor formation also remains an important element of concern with both plasma-derived and recombinant products, particularly with regard to FVIII in haemophilia A [87]. The risk of inhibitor formation was shown to be greater with recombinant versus plasma-derived factor VIII concentrates in some cohort studies [88], but similar in others [89].

In light of these considerations (availability, adverse reactions and cost), it appears that the issue of pathogen safety of blood/plasma-derived products is highly important but may not be the limiting factor with respect to overall patient safety. The benefits of treatment with a hypothetically 'unsafe' plasma-derived product may outweigh the risk of a negative outcome (e.g. bleeding, inhibitor formation), although we suggest that it may be clinically simpler to deal with inhibitor formation than to combat an infection from an unknown or untreatable pathogen.

7. Current knowledge gaps and areas of unmet need

There are still significant knowledge gaps and areas of unmet need with respect to the pathogen safety of blood/plasma-derived and recombinant products. The incidence of HBV, HCV and HIV TTI has fallen to near or below 1 per million transfused units in the industrialised world, indicating that current donor selection and blood screening strategies have had a positive impact on blood safety [14]. However, it is clear that screening both donors and donated blood cannot exclude all known pathogens or eliminate all risks

from emerging pathogens [63,90]. Historical precedent indicates that the blood supply is always vulnerable to contamination by hitherto non-prevalent/unknown pathogens, and that this risk cannot be accurately gauged [91]. As we identify new infectious agents of concern and develop new tests for their detection, it will also be necessary to clearly define the infectious dose range for each agent and use appropriately sensitive tests for their identification. For example, in suspected cases of vCJD infection, considerable challenges remain in the development of screening and confirmatory tests that have sufficient sensitivity and specificity to be of use in both a clinical setting and within blood banks [92].

Surveillance of people with haemophilia is required to monitor pathogen safety issues related to blood and plasma products. The European Haemophilia Safety Surveillance system (EUHASS), which began in 2008, is a pharmacovigilance programme which spans 25 European countries and is designed to detect, monitor and investigate adverse drug reactions. Reports of adverse events (such as acute/allergic events, TTIs and inhibitors) are submitted to EUHASS by participating centres and cumulative patient and clotting factor data are recorded annually [93].

A formal coordinated risk-assessment and management action plan, in addition to a task force, should be developed to respond quickly to any emerging infections. Such a plan should include long-term storage of samples from produced batches (for retesting in the case of outbreaks with known or recently emerged infectious agents) and guidance on responsibility for developing/performing tests for emerging pathogens (industry vs. regulatory agencies). Guidance on approaching patients who have potentially been infected and surveillance strategies for patients at high risk would also be beneficial.

8. Conclusions

The majority of evidence indicates that the concept of clinical safety of blood/plasma derivatives does not necessarily correspond to the concept of pathogen safety; blood can only be classed as microbially safe in reference to the infectious agents that are known and have been screened for. Establishing whether the presence of undetected microbes in the blood is clinically relevant will require further long-term, detailed studies. It should also be noted that even though the risk of transmission of key detectable viruses (such as HIV, HBV and HCV) via transfusion has fallen significantly, transmission does still occur.

In general, balancing safety, efficacy and practicalities is a difficult goal to achieve – patient safety is typically the key driver, but striving for near-complete safety at the expense of the patient's health or quality of life may not be the best course of action for patients or clinicians. The lack of a cohesive international strategy for blood donation and screening is a pressing concern that needs to be addressed. Furthermore, a formal coordinated continuous risk-assessment and management action plan needs to be developed to deal with the constant potential risk of emerging infections. Establishing an international registry (or harmonising data collection in National Registries) and dedicated task force may help to identify newly emerging pathogens more rapidly than in the past and to further improve pathogen safety of blood/plasma-derived products and the blood supply in general.

Practice points

- The use of blood/plasma-derived products for the treatment of bleeding disorders carries a risk of pathogen transmission.
- Blood donations are screened for key pathogens such as HBV, HCV, HIV and the causative agent of syphilis, but other screening tests should be conducted as required according to geographical location and patient risk factors.
- Screening tests for pathogens may lack sensitivity/specification and so false negatives may occur, resulting in a residual pathogen risk to patients.

- In terms of pathogen safety, recombinant products (products which have had minimal exposure to blood/plasma-derived proteins) are considered to pose the lowest level of risk to patients.

Research agenda

- Regional and international rates of transfusion-transmitted infection for key pathogens and emerging pathogens
- Safety and efficacy of blood/plasma-derived products and recombinant products for treatment of bleeding disorders.

Conflict of interest statement

A.T. has received grants and personal fees from Bayer, grants and personal fees from Baxter, grants and personal fees from Biostest, grants and personal fees from CSL Behring, grants and personal fees from Novo Nordisk, grants and personal fees from Pfizer, and grants and personal fees from Octapharma, during the conduct of the study.

C.F.P. has received grants as bureau speaker, consultant, or advisor, from Gilead, Merck Sharp and Dohme, Roche, Pfizer, Abbott, Bristol-Myers Squibb, VII, and Boehringer-Ingelheim. None of these personal activities is in conflict with the opinions he expressed in this manuscript.

D.N. has received honoraria for conferences from Pfizer, Roche Pharma, Roche Diagnostics, Abbott, MSD, and Astellas.

G.D.M. has disclosed the following financial relationships – speaker or a member of a speaker bureau for: Boehringer-Ingelheim, Sanofi-Aventis, Bayer, Novo Nordisk, Pfizer,

Biostest, and Grifols. Consultant or ad hoc speaker/consultant for: Boehringer-Ingelheim, Eli-Lilly, Sanofi-Aventis, Bayer, CSL Behring, Novo Nordisk, Pfizer, Biostest, and Grifols.

J.W.I. has received personal fees from Piramal and grants from the Department of Health, UK, outside the submitted work.

L.G. reports no potential conflicts of interest.

M.C. has received research grants, lecture fees, and honoraria for consultancy from Baxter, Bayer, and Pfizer.

Role of funding source

Funding for medical writing assistance and two author meetings were provided by Pfizer. Pfizer had no editorial influence regarding the contents of this manuscript and did not comment on the outline or any drafts prior to journal submission.

Acknowledgements

We wish to thank Professor Hermann Eichler for his appraisal of this manuscript and helpful suggestions. Medical writing assistance was provided by Hanna Mourad-Agha of inScience Communications, Springer Healthcare. This assistance was funded by Pfizer.

Appendix A. Supplementary data

Supplementary data to this article can be found online at <http://dx.doi.org/10.1016/j.blre.2015.07.004>.

References

- [1] Paessler S, Walker DH. Pathogenesis of the viral hemorrhagic fevers. *Annu Rev Pathol* 2013;8:411–40.
- [2] Srivastava A, Brewer AK, Mauser-Bunschoten EP, Key NS, Kitchen S, Llinas A, et al. Guidelines for the management of hemophilia. *Haemophilia* 2013;19:e1–47.
- [3] Benjamin RJ, McLaughlin LS. Plasma components: properties, differences, and uses. *Transfusion* 2012;52(Suppl. 1):9S–19S.
- [4] Stonebraker JS, Brooker M, Amand RE, Farrugia A, Srivastava A. A study of reported factor VIII use around the world. *Haemophilia* 2010;16:33–46.
- [5] Plug I, Van Der Bom JG, Peters M, Mauser-Bunschoten EP, De Goede-Bolder A, Heijnen L, et al. Mortality and causes of death in patients with hemophilia, 1992–2001: a prospective cohort study. *J Thromb Haemost* 2006;4:510–6.
- [6] Franchini M, Mannucci PM. Past, present and future of hemophilia: a narrative review. *Orphanet J Rare Dis* 2012;7:24.
- [7] Franchini M, Tagliaferri A, Mannucci PM. The management of hemophilia in elderly patients. *Clin Interv Aging* 2007;2:361–8.
- [8] Trimble SR, Parker CS, Grant AM, Soucie JM, Reyes N. Assessing emerging infectious threats to blood safety for the blood disorders community. *Am J Prev Med* 2010;38: S468–74.
- [9] Hardwick J. Blood processing. *ISBT Sci Ser* 2008;3:148–76.
- [10] World Health Organisation (WHO). WHO Technical Report: Guidelines on viral inactivation and removal procedures intended to assure the viral safety of human blood plasma products. 2004. Accessible from: http://www.who.int/bloodproducts/publications/WHO_TRS_924_A4.pdf.
- [11] Di Minno G, Canaro M, Ironside JW, Navarro D, Perno CF, Tiede A, et al. Pathogen safety of long-term treatments for bleeding disorders: still relevant to current practice. *Haematologica* 2013;98:1495–8.
- [12] Gringeri A. Factor VIII safety: plasma-derived versus recombinant products. *Blood Transfus* 2011;9:366–70.
- [13] Di Minno G, Canaro M, Ironside JW, Navarro D, Perno CF, Tiede A, et al. Pathogen safety of long-term treatments for bleeding disorders: (un)predictable risks and evolving threats. *Semin Thromb Hemost* 2013;39:973.
- [14] Stramer SL, Dodd RY. Transfusion-transmitted emerging infectious diseases: 30 years of challenges and progress. *Transfusion* 2013;53:2375–83.
- [15] Hourfar MK, Jork C, Schottstedt V, Weber-Schelle M, Brixner V, Busch MP, et al. Experience of German Red Cross blood donor services with nucleic acid testing: results of screening more than 30 million blood donations for human immunodeficiency virus-1, hepatitis C virus, and hepatitis B virus. *Transfusion* 2008;48:1558–66.
- [16] World Health Organisation (WHO). Recommendations: Screening Donated Blood for Transfusion-Transmissible Infections. 2009. Accessible from: <http://www.who.int/bloodsafety/ScreeningTTI.pdf>.
- [17] World Federation of Hemophilia (WFH). Registry of Clotting Factor Concentrates. 2012. Accessible from: <http://www1.wfh.org/publications/files/pdf-1227.pdf>.
- [18] Gerlich WH. Medical virology of hepatitis B: how it began and where we are now. *Virol J* 2013;10:239.
- [19] Ananworanich J, Fletcher JL, Pinyakorn S, van Grienden F, Vanderveen C, Schuetz A, et al. A novel acute HIV infection staging system based on 4th generation immunoassay. *Retrovirology* 2013;10:56.
- [20] Klamroth R, Groner A, Simon TL. Pathogen inactivation and removal methods for plasma-derived clotting factor concentrates. *Transfusion* 2014;54:1406–17.
- [21] Muller B, Nubling CM, Kress J, Roth WK, De Zolt S, Pichl L. How safe is safe: new human immunodeficiency virus Type 1 variants missed by nucleic acid testing. *Transfusion* 2013;53(Suppl. 3):2422–30.
- [22] Bruhn R, Lelie N, Custer B, Busch M, Kleinman S, International NATSG. Prevalence of human immunodeficiency virus RNA and antibody in first-time, lapsed, and repeat blood donations across five international regions and relative efficacy of alternative screening scenarios. *Transfusion* 2013;53:2399–412.
- [23] Datta S, Chatterjee S, Veer V. Recent advances in molecular diagnostics of hepatitis B virus. *World J Gastroenterol* 2014;20:14615–25.
- [24] Lieshout-Krikke RW, Molenaar-de Backer MW, van Swieten P, Zaaijer HL. Surface antigen-negative hepatitis B virus infection in Dutch blood donors. *Eur J Clin Microbiol* 2014;33:69–77.
- [25] Teo CG, Locarnini SA. Potential threat of drug-resistant and vaccine-escape HBV mutants to public health. *Antivir Ther* 2010;15:445–9.
- [26] Gupta E, Bajpai M, Choudhary A. Hepatitis C virus: Screening, diagnosis, and interpretation of laboratory assays. *Asian J Transfus Sci* 2014;8:19–25.
- [27] Kleinman S, Busch MP, Korelitz JJ, Schreiber GB. The incidence/window period model and its use to assess the risk of transfusion-transmitted human immunodeficiency virus and hepatitis C virus infection. *Transfus Med Rev* 1997;11:155–72.
- [28] Stramer SL, Krystof DE, Brodsky JP, Fickett TA, Reynolds B, Dodd RY, et al. Comparative analysis of triplex nucleic acid test assays in United States blood donors. *Transfusion* 2013;53(Suppl. 3):2525–37.
- [29] Sena AC, White BL, Sparling PF. Novel Treponema pallidum serologic tests: a paradigm shift in syphilis screening for the 21st century. *Clin Infect Dis* 2010; 51:700–8.
- [30] Pealer LN, Marfin AA, Petersen LR, Lanciotti RS, Page PL, Stramer SL, et al. Transmission of West Nile virus through blood transfusion in the United States in 2002. *N Engl J Med* 2003;349:1236–45.
- [31] Glynn SA, Busch MP, Dodd RY, Katz LM, Stramer SL, Klein HG, et al. Emerging infectious agents and the nation's blood supply: responding to potential threats in the 21st century. *Transfusion* 2013;53:438–54.
- [32] Pupella S, Pisani G, Cristiano K, Catalano L, Grazzini G. West Nile virus in the transfusion setting with a special focus on Italian preventive measures adopted in 2008–2012 and their impact on blood safety. *Blood Transfus* 2013;11:563–74.
- [33] Parola P, de Lamballerie X, Jourdan J, Rovery C, Vaillant V, Minodier P, et al. Novel chikungunya virus variant in travelers returning from Indian Ocean islands. *Emerg Infect Dis* 2006;12:1493–9.
- [34] Liument Bruno GM, Calteri D, Petropoulos K, Mattivi A, Po C, Macini P, et al. The Chikungunya epidemic in Italy and its repercussion on the blood system. *Blood Transfus* 2008;6:199–210.
- [35] Petersen LR, Epstein JS. Chikungunya virus: new risk to transfusion safety in the Americas. *Transfusion* 2014;54:1911–5.

- [36] Tsetsarkin KA, Vanlandingham DL, McGee CE, Higgs S. A single mutation in chikungunya virus affects vector specificity and epidemic potential. *PLoS Pathog* 2007;3:e201.
- [37] Norja P, Lassila R, Makris M. Parvovirus transmission by blood products - a cause for concern? *Br J Haematol* 2012;159:385–93.
- [38] Modrow S, Wenzel JJ, Schimanski S, Schwarzbek J, Rothe U, Oldenburg J, et al. Prevalence of nucleic acid sequences specific for human parvoviruses, hepatitis A and hepatitis E viruses in coagulation factor concentrates. *Vox Sang* 2011;100: 351–8.
- [39] Soucie JM, Monahan PE, Kulkarni R, De Staercke C, Recht M, Chitlur MB, et al. Evidence for the continued transmission of parvovirus B19 in patients with bleeding disorders treated with plasma-derived factor concentrates. *Transfusion* 2013;53: 1143–4.
- [40] Heegaard ED, Brown KE. Human parvovirus B19. *Clin Microbiol Rev* 2002;15: 485–505.
- [41] Ragni MV, Sherman KE, Jordan JA. Viral pathogens. *Haemophilia* 2010;16(Suppl. 5):40–6.
- [42] Pawlotsky JM. Hepatitis E screening for blood donations: an urgent need? *Lancet* 2014;384:1729–30.
- [43] Boxall E, Herborn A, Kochethu G, Pratt G, Adams D, Ijaz S, et al. Transfusion-transmitted hepatitis E in a 'nonhyperendemic' country. *Transfus Med* 2006;16:79–83.
- [44] Colson P, Coze C, Gallian P, Henry M, De Micco P, Tamalet C. Transfusion-associated hepatitis E, France. *Emerg Infect Dis* 2007;13:648–9.
- [45] Huzly D, Umhau M, Bettinger D, Cathomen T, Emmerich F, Hasselblatt P, et al. Transfusion-transmitted hepatitis E in Germany, 2013. *Euro Surveill* 2014;19.
- [46] Xu C, Wang RY, Schechter CA, Ge S, Shih JW, Xia NS, et al. An assessment of hepatitis E virus (HEV) in US blood donors and recipients: no detectable HEV RNA in 1939 donors tested and no evidence for HEV transmission to 362 prospectively followed recipients. *Transfusion* 2013;53(Suppl. 3):2505–11.
- [47] Dreier J, Juhl D. Autochthonous Hepatitis E Virus Infections: A New Transfusion-Associated Risk? *Transfus Med Hemother* 2014;41:29–39.
- [48] Takeda H, Matsubayashi K, Sakata H, Sato S, Kato T, Hino S, et al. A nationwide survey for prevalence of hepatitis E virus antibody in qualified blood donors in Japan. *Vox Sang* 2010;99:307–13.
- [49] Mansuy JM, Bendall R, Legrand-Abravanel F, Saune K, Miedouge M, Ellis V, et al. Hepatitis E virus antibodies in blood donors, France. *Emerg Infect Dis* 2011;17: 2309–12.
- [50] Hewitt PE, Ijaz S, Brailsford SR, Brett R, Dicks S, Haywood B, et al. Hepatitis E virus in blood components: a prevalence and transmission study in southeast England. *Lancet* 2014;384:1766–73.
- [51] Orru CD, Wilham JM, Vascellari S, Hughson AG, Caughey B. New generation QuIC assays for prion seeding activity. *Prion* 2012;6:147–52.
- [52] Morales R, Duran-Aniozzi C, Diaz-Espinoza R, Camacho MV, Soto C. Protein misfolding cyclic amplification of infectious prions. *Nat Protoc* 2012;7:1397–409.
- [53] Segarra C, Bougard D, Moudjou M, Laude H, Beringue V, Coste J. Plasminogen-based capture combined with amplification technology for the detection of PrP(TSE) in the pre-clinical phase of infection. *PLoS One* 2013;8:e69632.
- [54] Cooper JK, Andrews N, Ladhani K, Bujaki E, Minor PD. Evaluation of a test for its suitability in the diagnosis of variant Creutzfeldt-Jakob disease. *Vox Sang* 2013; 105:196–204.
- [55] Jackson GS, Burk-Rafel J, Edgeworth JA, Sicilia A, Abdilah S, Korteweg J, et al. Population Screening for Variant Creutzfeldt-Jakob Disease Using a Novel Blood Test: Diagnostic Accuracy and Feasibility Study. *JAMA Neurol* 2014;71:421–8.
- [56] Lacroux C, Comoy E, Moudjou M, Perret-Liaudet A, Lugan S, Litaise C, et al. Preclinical detection of variant CJD and BSE prions in blood. *PLoS Pathog* 2014;10: e1004202.
- [57] Irshad M, Joshi YK, Sharma Y, Dhar I. Transfusion transmitted virus: A review on its molecular characteristics and role in medicine. *World J Gastroenterol* 2006;12: 5122–34.
- [58] Stramer SL, Notari EP, Krysztof DE, Dodd RY. Hepatitis B virus testing by minipool nucleic acid testing: does it improve blood safety? *Transfusion* 2013;53(Suppl. 3):2449–58.
- [59] Salles NA, Levi JE, Barreto CC, Sampaio LP, Romano CM, Sabino EC, et al. Human immunodeficiency virus transfusion transmission despite nucleic acid testing. *Transfusion* 2013;53(Suppl. 3):2593–5.
- [60] Schmidt M, Korn K, Nubling CM, Chudy M, Kress J, Horst HA, et al. First transmission of human immunodeficiency virus Type 1 by a cellular blood product after mandatory nucleic acid screening in Germany. *Transfusion* 2009;49:1836–44.
- [61] Chudy M, Weber-Schehl M, Pichl L, Jork C, Kress J, Heiden M, et al. Blood screening nucleic acid amplification tests for human immunodeficiency virus Type 1 may require two different amplification targets. *Transfusion* 2012;52:431–9.
- [62] Paul-Ehrlich-Institut. Instruction to implement measures for risk minimisation in using HIV-1 NAT test systems. 2012. Available from: http://www.pei.de/SharedDocs/Downloads/vigilanz/haemovigilanz/beschreide/2012-06-15-implementation-hiv-1-nat-test-systems-dual-target.pdf?__blob=publicationFile&v=3.
- [63] Tsukada Y, Yokoyama K, Ishida A, Handa M, Mori T, Kizaki M, et al. Erythroid crisis caused by parvovirus B19 transmitted via red blood cell transfusion. *Intern Med* 2011;50:2379–82.
- [64] Lai L, Lee TH, Tobler L, Wen L, Shi P, Alexander J, et al. Relative distribution of West Nile virus RNA in blood compartments: implications for blood donor nucleic acid amplification technology screening. *Transfusion* 2012;52:447–54.
- [65] Goncalves DU, Proietti FA, Ribas JG, Araujo MG, Pinheiro SR, Guedes AC, et al. Epidemiology, treatment, and prevention of human T-cell leukemia virus type 1-associated diseases. *Clin Microbiol Rev* 2010;23:577–89.
- [66] Lindholm PF, Annen K, Ramsey G. Approaches to minimize infection risk in blood banking and transfusion practice. *Infect Disord Drug Targets* 2011;11:45–56.
- [67] Yesli SajAO. Minimum Infective Dose of the Major Human Respiratory and Enteric Viruses Transmitted Through Food and the Environment. *Food Environ Virol* 2011; 3:1–30.
- [68] Leggett HC, Cornwallis CK, West SA. Mechanisms of pathogenesis, infective dose and virulence in human parasites. *PLoS Pathol* 2012;8:e1002512.
- [69] Boeckh M, Geballe AP. Cytomegalovirus: pathogen, paradigm, and puzzle. *J Clin Invest* 2011;121:1673–80.
- [70] Souza MA, Passos AM, Treitinger A, Spada C. Seroprevalence of cytomegalovirus antibodies in blood donors in southern, Brazil. *Rev Soc Bras Med Trop* 2010;43: 359–61.
- [71] Ringwald J, Mertz I, Zimmermann R, Weisbach V, Strasser E, Achenbach S, et al. Hepatitis B virus vaccination of blood donors—what costs may be expected? *Transfus Med* 2005;15:83–92.
- [72] Fischinger JM, Stephan B, Wasserscheid K, Eichler H, Gartner BC. A cost-benefit analysis of blood donor vaccination as an alternative to additional DNA testing for reducing transfusion transmission of hepatitis B virus. *Vaccine* 2010;28: 7797–802.
- [73] World Health Organisation (WHO). WHO Technical Report, Series No. 924, 2004. Guidelines on viral inactivation and removal procedures intended to assure the viral safety of human blood plasma products. Accessible from: http://www.who.int/bloodproducts/publications/WHO_TRS_924_A4.pdf.
- [74] Roth WK. Quarantine Plasma: Quo vadis? *Transfus Med Hemother* 2010;37: 118–22.
- [75] Klein HG. Pathogen inactivation technology: cleansing the blood supply. *J Intern Med* 2005;257:224–37.
- [76] Schneider B, Becker M, Brackmann HH, Eis-Hubinger AM. Contamination of coagulation factor concentrates with human parvovirus B19 genotype 1 and 2. *Thromb Haemost* 2004;92:838–45.
- [77] Roberts PL, Dalton J, Evans D, Harrison P, Li Z, Ternouth K, et al. Removal of TSE agent from plasma products manufactured in the United Kingdom. *Vox Sang* 2013;104:299–308.
- [78] Cai K, Osheroff WP, Buczynski G, Hotta J, Lang J, Elliott E, et al. Characterization of Thrombate III(R), a pasteurized and nanofiltered therapeutic human antithrombin concentrate. *Biologicals* 2014;42:133–8.
- [79] Cardone F, Simoneau S, Arzel A, Puopolo M, Berardi VA, Abdel-Haq H, et al. Comparison of nanofiltration efficacy in reducing infectivity of centrifuged versus ultracentrifuged 263 K scrapie-infected brain homogenates in "spiked" albumin solutions. *Transfusion* 2012;52:953–62.
- [80] Marschner S, Goodrich R. Pathogen Reduction Technology Treatment of Platelets, Plasma and Whole Blood Using Riboflavin and UV Light. *Transfus Med Hemother* 2011;38:8–18.
- [81] Seltsam A, Muller TH. UVC Irradiation for Pathogen Reduction of Platelet Concentrates and Plasma. *Transfus Med Hemother* 2011;38:43–54.
- [82] Reddy HL, Doane SK, Keil SD, Marschner S, Goodrich RP. Development of a riboflavin and ultraviolet light-based device to treat whole blood. *Transfusion* 2013; 53(Suppl. 1):131S–63.
- [83] Keeling D, Tait C, Makris M. Guideline on the selection and use of therapeutic products to treat haemophilia and other hereditary bleeding disorders. A United Kingdom Haemophilia Center Doctors' Organisation (UKHCDO) guideline approved by the British Committee for Standards in Haematology. *Haemophilia* 2008;14:671–84.
- [84] Kato H, Uruma M, Okuyama Y, Fujita H, Handa M, Tomiyama Y, et al. Incidence of transfusion-related adverse reactions per patient reflects the potential risk of transfusion therapy in Japan. *Am J Clin Pathol* 2013;140:219–24.
- [85] Directive 2002/98/EC of the European Parliament and of the Council of 27 January 2003 setting standards of quality and safety for the collection, testing, processing, storage and distribution of human blood and blood components and amending Directive 2001/83/EC. Available from: <http://eur-lex.europa.eu/legal-content/EN/TXT/?uri=CELEX:32002L0098>. [accessed Feb 2015].
- [86] Grillberger L, Kreil TR, Nasr S, Reiter M. Emerging trends in plasma-free manufacturing of recombinant protein therapeutics expressed in mammalian cells. *Biotechnol J* 2009;4:186–201.
- [87] AleDort LM. Is the incidence and prevalence of inhibitors greater with recombinant products? Yes. *J Thromb Haemost* 2004;2:861–2.
- [88] Goudemand J, Rothchild C, Demiguel V, Vinciguerra C, Lambert T, Chambost H, et al. Influence of the type of factor VIII concentrate on the incidence of factor VIII inhibitors in previously untreated patients with severe hemophilia A. *Blood* 2006;107:46–51.
- [89] Gouw SC, van den Berg HM, Fischer K, Auerswald G, Carcao M, Chalmers E, et al. Intensity of factor VIII treatment and inhibitor development in children with severe hemophilia A: the RODIN study. *Blood* 2013;121:4046–55.
- [90] Ludlam CA, Powderly WG, Bozzette S, Diamond M, Koerner MA, Kulkarni R, et al. Clinical perspectives of emerging pathogens in bleeding disorders. *Lancet* 2006; 367:252–61.
- [91] Knight R, Stanley S, Wong M, Dolan G. Hemophilia therapy and blood-borne pathogen risk. *Semin Thromb Hemost* 2006;32(Suppl. 2):3–9.
- [92] Brown P. Creutzfeldt-Jakob disease: reflections on the risk from blood product therapy. *Haemophilia* 2007;13(Suppl. 5):33–40.
- [93] Makris M, Calizzani G, Fischer K, Gilman EA, Hay CR, Lassila R, et al. EUHASS: The European Haemophilia Safety Surveillance system. *Thromb Res* 2011;127(Suppl. 2):S22–5.
- [94] Peden A, McCordle L, Head MW, Love S, Ward HJ, Cousens SN, et al. Variant CJD infection in the spleen of a neurologically asymptomatic UK adult patient with hemophilia. *Haemophilia* 2010;16:296–304.
- [95] Edgeworth JA, Farmer M, Sicilia A, Tavares P, Beck J, Campbell T, et al. Detection of prion infection in variant Creutzfeldt-Jakob disease: a blood-based assay. *Lancet* 2011;377:487–93.

- [96] World Health Organisation (WHO). Factsheet No 327 - Chikungunya. 2014. Accessible from: <http://www.who.int/mediacentre/factsheets/fs327/en/>.
- [97] Laurent P, Le Roux K, Grivard P, Bertil G, Naze F, Picard M, et al. Development of a sensitive real-time reverse transcriptase PCR assay with an internal control to detect and quantify chikungunya virus. *Clin Chem* 2007;53:1408–14.
- [98] Santhosh SR, Parida MM, Dash PK, Pateriya A, Pattnaik B, Pradhan HK, et al. Development and evaluation of SYBR Green I-based one-step real-time RT-PCR assay for detection and quantification of Chikungunya virus. *J Clin Virol* 2007;39:188–93.
- [99] Blacksell SD, Tanganuchitcharnchai A, Jarman RG, Gibbons RV, Paris DH, Bailey MS, et al. Poor diagnostic accuracy of commercial antibody-based assays for the diagnosis of acute Chikungunya infection. *Clin Vaccin Immunol* 2011;18:1773–5.
- [100] CMV HHV-6,7,8 R-gene® Real-Time PCR Kit Technical Specifications. 2015. Available from: <http://www.biomerieux-diagnostics.com/cmv-hhv678-r-gener>.
- [101] Michelin BD, Hadzisejdic I, Bozic M, Grahovac M, Hess M, Grahovac B, et al. Detection of cytomegalovirus (CMV) DNA in EDTA whole-blood samples: evaluation of the quantitative artus CMV LightCycler PCR kit in conjunction with automated sample preparation. *J Clin Microbiol* 2008;46:1241–5.
- [102] Teoh BT, Sam SS, Tan KK, Johari J, Danlami MB, Hooi PS, et al. Detection of dengue viruses using reverse transcription-loop-mediated isothermal amplification. *BMC Infect Dis* 2013;13:387.
- [103] Simplexa™ Dengue Kit Technical Specifications. 2015. Available from: <http://www.focusdx.com/product/MOL300/ous>.
- [104] Chao DY, Davis BS, Chang GJ. Development of multiplex real-time reverse transcriptase PCR assays for detecting eight medically important flaviviruses in mosquitoes. *J Clin Microbiol* 2007;45:584–9.
- [105] Hunsperger EA, Yoksan S, Buchy P, Nguyen VC, Sekaran SD, Enria DA, et al. Evaluation of commercially available diagnostic tests for the detection of dengue virus NS1 antigen and anti-dengue virus IgM antibody. *PLoS Negl Trop Dis* 2014;8:e3171.
- [106] World Health Organisation (WHO). Factsheet No 360 - HIV/AIDS. Accessible from: <http://www.who.int/mediacentre/factsheets/fs360/en/>.
- [107] Procleix® UltraTrio® Assay. 2015. Available from: <http://www.fda.gov/ucm/groups/fdagov-public/@fdagov-bio-gen/documents/document/ucm335285.pdf>.
- [108] Shah SMSS. Hepatitis B virus serology: Use and interpretation. *Hep B Annual* 2007; 4:39–54.
- [109] Ba Alawi F, Robertson PW, LePage AK, Jayamaha J, Baleriola C, Rawlinson WD. The reliability of HBV core antibody in serological screening for hepatitis B virus. *Pathology* 2013;45:501–5.
- [110] World Health Organisation (WHO). Factsheet No 164 - Hepatitis C. 2013. Accessible from: <http://www.who.int/mediacentre/factsheets/fs164/en/>.
- [111] NGI UltraQualITM HCV RT-PCR assay. 2015. Available from: <http://www.fda.gov/downloads/biologicsbloodvaccines/ucm148271.pdf>.
- [112] Kamili S, Drobniuc J, Araujo AC, Hayden TM. Laboratory diagnostics for hepatitis C virus infection. *Clin Infect Dis* 2012;55(Suppl. 1):S43–8.
- [113] Tillmann HL. Hepatitis C virus core antigen testing: role in diagnosis, disease monitoring and treatment. *World J Gastroenterol* 2014;20:6701–6.
- [114] van Helden J, Weiskirchen R. Hepatitis C diagnostics: clinical evaluation of the HCV-core antigen determination. *Z Gastroenterol* 2014;52:1164–70.
- [115] Bajpai M, Gupta E. Transfusion-transmitted hepatitis E: is screening warranted? *Indian J Med Microbiol* 2011;29:353–8.
- [116] Vollmer T, Knabbe C, Dreier J. Comparison of real-time PCR and antigen assays for detection of hepatitis E virus in blood donors. *J Clin Microbiol* 2014;52: 2150–6.
- [117] Abravanel F, Sandres-Saune K, Lhomme S, Dubois M, Mansuy JM, Izopet J. Genotype 3 diversity and quantification of hepatitis E virus RNA. *J Clin Microbiol* 2012;50: 897–902.
- [118] Vollmer T, Diekmann J, Johnne R, Eberhardt M, Knabbe C, Dreier J. Novel approach for detection of hepatitis E virus infection in German blood donors. *J Clin Microbiol* 2012;50:2708–13.
- [119] Drobniuc J, Meng J, Reuter G, Greene-Montfort T, Khudyakova N, Dimitrova Z, et al. Serologic assays specific to immunoglobulin M antibodies against hepatitis E virus: pangenotypic evaluation of performances. *Clin Infect Dis* 2010;51: e24–7.
- [120] Henke-Gendo C, Schulz TF. Transmission and disease association of Kaposi's sarcoma-associated herpesvirus: recent developments. *Curr Opin Infect Dis* 2004;17: 53–7.
- [121] HHV-8 Quantitative Real-time PCR. 2015. Available from: <http://www.viracoribt.com/Test-Catalog/Detail/Human-Herpes-Virus-8-HHV-8-Quantitative-PCR-8000>.
- [122] Mbisa GL, Miley W, Gamache CJ, Gillette WK, Esposito D, Hopkins R, et al. Detection of antibodies to Kaposi's sarcoma-associated herpesvirus: a new approach using K8.1 ELISA and a newly developed recombinant LANA ELISA. *J Immunol Methods* 2010;356:39–46.
- [123] United Nations (UN). Global report: UNAIDS report on the global AIDS epidemic. 2013. Accessible from <http://www.unaids.org/en/resources/publications/2013/name,85053en.asp>.
- [124] Naeth G, Ehret R, Wiesmann F, Braun P, Knechten H, Berger A. Comparison of HIV-1 viral load assay performance in immunological stable patients with low or undetectable viremia. *Med Microbiol Immunol* 2013;202:67–75.
- [125] Butto S, Suligoi B, Fanales-Belasio E, Raimondo M. Laboratory diagnostics for HIV infection. *Ann Ist Super Sanita* 2010;46:24–33.
- [126] Lairmore MD, Anupam R, Bowden N, Haines R, Haynes 2nd RA, Ratner L, et al. Molecular determinants of human T-lymphotropic virus type 1 transmission and spread. *Viruses* 2011;3:1131–65.
- [127] Abbott Prism Human T-Lymphotropic Virus Types I and II assay. Available from: <http://www.fda.gov/downloads/BiologicsBloodVaccines/BloodBloodProducts/ApprovedProducts/LicensedProductsBLAs/BloodDonorScreening/InfectiousDisease/UCM298517.pdf>.
- [128] Avoq HTLV-I/II Microelisa system. Available from: <http://www.fda.gov/downloads/BiologicsBloodVaccines/BloodBloodProducts/ApprovedProducts/LicensedProductsBLAs/BloodDonorScreening/InfectiousDisease/UCM298517.pdf>.
- [129] Rocha-Junior MH R, Wagatsuma V, Takayanagi O, Slavov S, Otaguri K, Rodrigues E, et al. Development and validation of a Multiplex Real-Time PCR for HTLV-1/2 confirmatory diagnosis. *Retrovirology* 2014;11:105.
- [130] Malm K, Kjerstadius T, Andersson S. Evaluation of a new screening assay for HTLV-1 and -2 antibodies for large-scale use. *J Med Virol* 2010;82:1606–11.
- [131] Andrade RG, Ribeiro MA, Namen-Lopes MS, Silva SM, Basques FV, Ribas JG, et al. Evaluation of the use of real-time PCR for human T cell lymphotropic virus 1 and 2 as a confirmatory test in screening for blood donors. *Rev Soc Bras Med Trop* 2010;43:111–5.
- [132] Bonvicini F, Manaresi E, Bua G, Venturoli S, Gallinella G. Keeping pace with Parvovirus B19 Genetic Variability: a Multiplex Genotype-Specific qPCR Assay. *J Clin Microbiol* 2013.
- [133] Koppelman MH, Cuypers HT, Emrich T, Zaaijer HL. Quantitative real-time detection of parvovirus B19 DNA in plasma. *Transfusion* 2004;44:97–103.
- [134] Tong R, Zhou WM, Liu XJ, Wang Y, Lou YL, Tan WJ. Detection of human parvovirus B19, human bocavirus and human parvovirus 4 infections in blood samples among 95 patients with liver disease in Nanjing by nested PCR. *Zhonghua Shi Yan He Lin Chuang Bing Du Xue Za Zhi* 2013;27:135–7.
- [135] Doyle S, Kerr S, O'Keeffe G, O'Carroll D, Daly P, Kilty C. Detection of parvovirus B19 IgM by antibody capture enzyme immunoassay: receiver operating characteristic analysis. *J Virol Methods* 2000;90:143–52.
- [136] Chen MY, Yang SJ, Hung CC. Placental transmission of human parvovirus 4 in newborns with hydrops, Taiwan. *Emerg Infect Dis* 2011;17:1954–6.
- [137] Lahtinen A, Kivelä P, Hedman I, Kumar A, Kantele A, Lappalainen M, et al. Serodiagnosis of primary infections with human parvovirus 4, Finland. *Emerg Infect Dis* 2011;17:79–82.
- [138] Lavoie M, Sharp CP, Pepin J, Pennington C, Fouppouapouognigni Y, Pybus OG, et al. Human parvovirus 4 infection, Cameroon. *Emerg Infect Dis* 2012;18:680–3.
- [139] May J, Drexler JF, Reber U, Sarpong N, Adjei O, Panning M, et al. Human parvovirus 4 viremia in young children, Ghana. *Emerg Infect Dis* 2012;18:1690–2.
- [140] Fryer JF, Kapoor A, Minor PD, Delwart E, Baylis SA. Novel parvovirus and related variant in human plasma. *Emerg Infect Dis* 2006;12:151–4.
- [141] Simmonds P, Davidson F, Lycett C, Prescott LE, MacDonald DM, Ellender J, et al. Detection of a novel DNA virus (TTV) in blood donors and blood products. *Lancet* 1998;352:191–5.
- [142] Tyagi AK, Pradier A, Baumer O, Uppugunduri CR, Huezo-Diaz P, Posfay-Barbe KM, et al. Validation of SYBR Green based quantification assay for the detection of human Torque Teno virus titers from plasma. *Virology* 2013;10:191.
- [143] West Nile Virus: Detection with Serologic and Real-time PCR Assays. Quest Diagnostics. 2015. Available from: http://www.questdiagnostics.com/testcenter/testguide.action?dc=CF_WestNileVirus.
- [144] Sambri V, Capobianchi MR, Cavrini F, Charrel R, Donoso-Mantke O, Escadafal C, et al. Diagnosis of west nile virus human infections: overview and proposal of diagnostic protocols considering the results of external quality assessment studies. *Viruses* 2013;5:2329–48.
- [145] Sanchini A, Donoso-Mantke O, Papa A, Sambri V, Teichmann A, Niedrig M. Second international diagnostic accuracy study for the serological detection of West Nile virus infection. *PLoS Negl Trop Dis* 2013;7:e2184.
- [146] World Health Organisation (WHO). Global incidence and prevalence of selected curable sexually transmitted infections. 2008. Accessible from: http://apps.who.int/iris/bitstream/10665/75181/1/9789241503839_eng.pdf.
- [147] Palmer HM, Higgins SP, Herring AJ, Kingston MA. Use of PCR in the diagnosis of early syphilis in the United Kingdom. *Sex Transm Infect* 2003;79:479–83.
- [148] Grange PA, Gressier L, Dion PL, Farhi D, Benhaddou N, Gerhardt P, et al. Evaluation of a PCR test for detection of treponema pallidum in swabs and blood. *J Clin Microbiol* 2012;50:546–52.
- [149] Ratnam S. The laboratory diagnosis of syphilis. *Can J Infect Dis Med Microbiol* 2005; 16:45–51.
- [150] Mosqueda J, Olvera-Ramirez A, Aguilar-Tipacamú G, Canto GJ. Current advances in detection and treatment of babesiosis. *Curr Med Chem* 2012;19:1504–18.
- [151] Teal AE, Habura A, Ennis J, Keithly JS, Madison-Antenucci S. A new real-time PCR assay for improved detection of the parasite Babesia microti. *J Clin Microbiol* 2012;50:903–8.
- [152] Priest JW, Moss DM, Won K, Todd CW, Henderson L, Jones CC, et al. Multiplex assay detection of immunoglobulin G antibodies that recognize Babesia microti antigens. *Clin Vaccine Immunol* 2012;19:1539–48.
- [153] World Health Organisation (WHO). Factsheet No 375 - Leishmaniasis. 2014. Accessible from: <http://www.who.int/mediacentre/factsheets/fs375/en/>.
- [154] Kobets T, Grekov I, Lipordova M. Leishmaniasis: prevention, parasite detection and treatment. *Curr Med Chem* 2012;19:1443–74.
- [155] Abbasi I, Aramin S, Hailu A, Shiferaw W, Kassahun A, Belay S, et al. Evaluation of PCR procedures for detecting and quantifying Leishmania donovani DNA in large numbers of dried human blood samples from a visceral leishmaniasis focus in Northern Ethiopia. *BMC Infect Dis* 2013;13:153.
- [156] Bensoussan E, Nasereddin A, Jonas F, Schnur LF, Jaffe CL. Comparison of PCR assays for diagnosis of cutaneous leishmaniasis. *J Clin Microbiol* 2006;44:1435–9.
- [157] World Health Organisation (WHO). Factsheet No 94 - Malaria. 2014. Accessible from: <http://www.who.int/mediacentre/factsheets/fs094/en/>.
- [158] Cordray MS, Richards-Kortum RR. Emerging nucleic acid-based tests for point-of-care detection of malaria. *Am J Trop Med Hyg* 2012;87:223–30.

- [159] Rodrigues MH, Cunha MG, Machado RL, Ferreira Jr OC, Rodrigues MM, Soares IS. Serological detection of *Plasmodium vivax* malaria using recombinant proteins corresponding to the 19-kDa C-terminal region of the merozoite surface protein-1. *Malar J* 2003;2:39.
- [160] World Health Organisation (WHO). Factsheet No 259 -Trypanosomiasis. 2014. Accessible from: <http://www.who.int/mediacentre/factsheets/fs259/en/>.
- [161] Chappuis F, Loutan L, Simarro P, Lejon V, Buscher P. Options for field diagnosis of human african trypanosomiasis. *Clin Microbiol Rev* 2005;18:133–46.
- [162] Mugasa CM, Laurent T, Schoone GJ, Kager PA, Lubega GW, Schallig HD. Nucleic acid sequence-based amplification with oligochromatography for detection of *Trypanosoma brucei* in clinical samples. *J Clin Microbiol* 2009;47:630–5.
- [163] Deborggraeve S, Lejon V, Ekangu RA, Mumba Ngoyi D, Pati Pyana P, Ilunga M, et al. Diagnostic accuracy of PCR in gambiense sleeping sickness diagnosis, staging and post-treatment follow-up: a 2-year longitudinal study. *PLoS Negl Trop Dis* 2011;5:e972.
- [164] World Health Organisation (WHO). Factsheet No 340 - Chagas Disease. 2014. Accessible from: <http://www.who.int/mediacentre/factsheets/fs340/en/>.
- [165] Schijman AG, Bisio M, Orellana L, Sued M, Duffy T, Mejia Jaramillo AM, et al. International study to evaluate PCR methods for detection of *Trypanosoma cruzi* DNA in blood samples from Chagas disease patients. *PLoS Negl Trop Dis* 2011;5:e931.
- [166] Caballero ZC, Sousa OE, Marques WP, Saez-Alquezar A, Umezawa ES. Evaluation of serological tests to identify *Trypanosoma cruzi* infection in humans and determine cross-reactivity with *Trypanosoma rangeli* and *Leishmania* spp. *Clin Vaccine Immunol* 2007;14:1045–9.
- [167] Grard G, Fair JN, Lee D, Slikas E, Steffen I, Muyembe JJ, et al. A novel rhabdovirus associated with acute hemorrhagic fever in central Africa. *PLoS Pathog* 2012;8:e1002924.
- [168] Bao CJ, Qi X, Wang H. A novel bunyavirus in China. *N Engl J Med* 2011;365:862–3 [author reply 864–865].
- [169] Zhang YZ, He YW, Dai YA, Xiong Y, Zheng H, Zhou DJ, et al. Hemorrhagic fever caused by a novel Bunyavirus in China: pathogenesis and correlates of fatal outcome. *Clin Infect Dis* 2012;54:527–33.
- [170] Christian KA, Ijaz K, Dowell SF, Chow CC, Chitale RA, Bresee JS, et al. What we are watching—five top global infectious disease threats, 2012: a perspective from CDC's Global Disease Detection Operations Center. *Emerg Health Threats J* 2013;6:20632.
- [171] El-Zoghby EF, Aly MM, Nasef SA, Hassan MK, Arafa AS, Selim AA, et al. Surveillance on A/H5N1 virus in domestic poultry and wild birds in Egypt. *Virol J* 2013;10:203.
- [172] Cowling BJ, Jin L, Lau EH, Liao Q, Wu P, Jiang H, et al. Comparative epidemiology of human infections with avian influenza A H7N9 and H5N1 viruses in China: a population-based study of laboratory-confirmed cases. *Lancet* 2013;382:129–37.
- [173] To KK, Chan JF, Yuen KY. Viral lung infections: Epidemiology, virology, clinical features, and management of avian influenza A(H7N9). *Curr Opin Pulm Med* 2014;20:225–32.
- [174] Briese T, Paweska JT, McMullan LK, Hutchison SK, Street C, Palacios G, et al. Genetic detection and characterization of Lujo virus, a new hemorrhagic fever-associated arenavirus from southern Africa. *PLoS Pathog* 2009;5:e1000455.
- [175] Paweska JT, Sewlall NH, Ksiazek TG, Blumberg LH, Hale MJ, Lipkin WI, et al. Nosocomial outbreak of novel arenavirus infection, southern Africa. *Emerg Infect Dis* 2009;15:1598–602.
- [176] Popgeorgiev N, Boyer M, Fancello L, Monteil S, Robert C, Rivet R, et al. Marseillevirus-like virus recovered from blood donated by asymptomatic humans. *J Infect Dis* 2013;208:1042–50.
- [177] Popgeorgiev N, Colson P, Thuret I, Chiarioni J, Gallian P, Raoult D, et al. Marseillevirus prevalence in multitransfused patients suggests blood transmission. *J Clin Virol* 2013;58:722–5.
- [178] Cotten M, Watson SJ, Kellam P, Al-Rabeeah AA, Makhdoom HQ, Assiri A, et al. Transmission and evolution of the Middle East respiratory syndrome coronavirus in Saudi Arabia: a descriptive genomic study. *Lancet* 2013.
- [179] Breban R, Riou J, Fontanet A. Interhuman transmissibility of Middle East respiratory syndrome coronavirus: estimation of pandemic risk. *Lancet* 2013;382:694–9.
- [180] Centers for Disease Control and Prevention (CDC). Severe Acute Respiratory Syndrome (SARS) – frequently asked questions. Accessible from: <http://www.cdc.gov/sars/about/faq.html>.
- [181] Nie QH, Luo XD, Zhang JZ, Su Q. Current status of severe acute respiratory syndrome in China. *World J Gastroenterol* 2003;9:1635–45.
- [182] Peiris JS, Guan Y. Confronting SARS: a view from Hong Kong. *Philos Trans R Soc Lond B Biol Sci* 2004;359:1075–9.
- [183] Mishra N, Pereira M, Rhodes RH, An P, Pipas JM, Jain K, et al. Identification of a novel polyomavirus in a pancreatic transplant recipient with retinal blindness and vasculitic myopathy. *J Infect Dis* 2014;210:1595–9.
- [184] Sauvage V, Livartowski A, Boizeau L, Servant-Delmas A, Lionnet F, Lefrere JJ, et al. No evidence of Marseillevirus-like virus presence in blood donors and recipients of multiple blood transfusions. *J Infect Dis* 2014;210:2017–8.
- [185] Cohen MS, Gay CL, Busch MP, Hecht FM. The detection of acute HIV infection. *J Infect Dis* 2010;202(Suppl. 2):S270–7.
- [186] Ciccaglione AR, Miceli M, Pisani G, Bruni R, Iudicone P, Costantino A, et al. Improving HIV-2 detection by a combination of serological and nucleic acid amplification test assays. *J Clin Microbiol* 2010;48:2902–8.
- [187] Food and Drug Administration (FDA) Guidance for Industry: Use of Nucleic Acid Tests on Pooled and Individual Samples from Donors of Whole Blood and Blood Components, Including Source Plasma, to Reduce the Risk of Transmission of Hepatitis B Virus. Accessible from: <http://www.fda.gov/BiologicsBloodVaccines/GuidanceComplianceRegulatoryInformation/Guidances/Blood/ucm327850.htm>; Jan 2015.
- [188] Namululi BA, Guerrieri C, Dramaix MW. Prevalence and incidence of HIV and hepatitis B among blood donors and estimated residual risk of transmission of HIV and HBV virus by blood transfusion. A study at the Provincial General Referee Hospital Bukavu, Democratic Republic of the Congo. *Rev Epidemiol Sante Publique* 2013;61:139–44.
- [189] de Almeida-Neto C, Goncalez TT, Birch RJ, de Carvalho SM, Capuani L, Leao SC, et al. Risk factors for human immunodeficiency virus infection among Brazilian blood donors: a multicentre case-control study using audio computer-assisted structured interviews. *Vox Sang* 2013;105:91–9.
- [190] Rosenberg GK, Lattimore S, Brailsford SR, Hewitt PE, Tettmar KJ, Kitchen AD, et al. The diversity of chronic hepatitis B virus infections within blood donors in England and North Wales 2005 through 2010. *Transfusion* 2013;53:2467–76.
- [191] Seed CR, Kiely P. A method for estimating the residual risk of transfusion-transmitted HBV infection associated with occult hepatitis B virus infection in a donor population without universal anti-HBc screening. *Vox Sang* 2013;105:290–8.
- [192] Offergeld R, Ritter S, Hamouda O. [HIV, HCV, HBV and syphilis surveillance among blood donors in Germany 2008–2010]. *Bundesgesundheitsblatt, Gesundheitsforschung, Gesundheitsschutz* 2012;55:907–13.
- [193] Koch C, Araujo F. Evolution of residual risk for HIV, HCV and HBV, from 1999 to 2010, in blood donations of the Centro Hospitalar S. Joao, EPE, Porto, Portugal. *Acta Med Port* 2013;26:371–6.
- [194] Cleland A, Smith L, Crossan C, Blatchford O, Dalton HR, Scobie L, et al. Hepatitis E virus in Scottish blood donors. *Vox Sang* 2013;105:283–9.
- [195] Baylis SA, Gartner T, Nick S, Ovemyr J, Blumel J. Occurrence of hepatitis E virus RNA in plasma donations from Sweden, Germany and the United States. *Vox Sang* 2012;103:89–90.
- [196] World Federation of Hemophilia (WFH) Annual Global Surveys. Accessible from: <http://www.wfh.org/en/page.aspx?pid=878>.