




Eptacog beta efficacy and safety in the treatment and control of bleeding in paediatric subjects (<12 years) with haemophilia A or B with inhibitors

Steven W. Pipe¹  | Cédric Hermans²  | Meera Chitlur³ | Manuel Carcao⁴  |
 Giancarlo Castaman⁵  | Joanna A. Davis⁶ | Jonathan Ducore⁷ | Amy L. Dunn⁸  |
 Miguel Escobar⁹  | Janna Journeycake¹⁰ | Osman Khan¹⁰ | Johnny Mahlangu¹¹  |
 Shannon L. Meeks¹²  | Ismail Haroon Mitha¹³ | Claude Négrier¹⁴ |
 Ulrike Nowak-Göttl¹⁵ | Michael Recht^{16,17} | Tammuela Chrisentery-Singleton¹⁸ |
 Oleksandra Stasyshyn¹⁹ | Kateryna V. Vilchevska²⁰ | Laura Villarreal Martinez²¹  |
 Michael Wang²²  | Jerzy Windyga²³ | Guy Young^{24,25}  | W. Allan Alexander²⁶ |
 Daniel Bonzo²⁷ | Christopher Macie²⁸ | Ian S. Mitchell²⁸ | Evelyne Sauty²⁹ |
 Thomas A. Wilkinson³⁰ | Amy D. Shapiro³¹

¹University of Michigan, Ann Arbor, Michigan, USA

²Cliniques Saint-Luc, Université Catholique de Louvain, Brussels, Belgium

³Children's Hospital of Michigan, Central Michigan University, Detroit, Michigan, USA

⁴The Hospital for Sick Children, Toronto, Ontario, Canada

⁵Center for Bleeding Disorders and Coagulation, Careggi University Hospital, Florence, Italy

⁶Pediatric Hemophilia Treatment Center, University of Miami, Miami, Florida, USA

⁷Hematology/Oncology Clinic, University of California at Davis, Sacramento, California, USA

⁸Nationwide Children's Hospital, Department of Pediatrics at The Ohio State University College of Medicine, Columbus, Ohio, USA

⁹University of Texas Health Science Center at Houston, Houston, Texas, USA

¹⁰Oklahoma Center for Bleeding and Clotting Disorders at OU Health, Oklahoma City, Oklahoma, USA

¹¹Hemophilia Comprehensive Care Center, University of the Witwatersrand and National Health Laboratory Service, Johannesburg, South Africa

¹²Emory University and Aflac Cancer and Blood Disorders Center of Children's Healthcare of Atlanta, Atlanta, Georgia, USA

¹³Lakeview Hospital, Benoni, Gauteng, South Africa

¹⁴Edouard Herriot University Hospital, Lyon, France

¹⁵Institute of Clinical Chemistry, University Hospital Schleswig-Holstein, Kiel, Germany

¹⁶American Thrombosis and Hemostasis Network, Rochester, New York, USA

¹⁷Oregon Health & Science University, Portland, Oregon, USA

¹⁸Louisiana Center for Advanced Medicine, Slidell, Louisiana, USA

¹⁹Institute of Blood Pathology and Transfusion Medicine, Lviv, Ukraine

²⁰National Specialized Children's Hospital Okhmatdyt, Kyiv, Ukraine

²¹Dr. José Eleuterio González Monterrey University Hospital, Monterrey, Nuevo León, México

²²Hemophilia and Thrombosis Center, University of Colorado, Aurora, Colorado, USA

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2022 The Authors. *Haemophilia* published by John Wiley & Sons Ltd.

²³Department of Hemostasis Disorders and Internal Medicine, Institute of Hematology and Transfusion Medicine, Warsaw, Poland

²⁴Children's Hospital Los Angeles, Los Angeles, California, USA

²⁵Keck School of Medicine, University of Southern California, Los Angeles, California, USA

²⁶Aoede Associates, Athens, Texas, USA

²⁷LFB-USA, Inc., Framingham, Massachusetts, USA

²⁸HEMA Biologics, LLC, Louisville, Kentucky, USA

²⁹LFB, Laboratoire français du fractionnement et des biotechnologies, Les Ulis, France

³⁰GLOVAL LLC, Broomfield, Colorado, USA

³¹Indiana Hemophilia and Thrombosis Center, Indianapolis, Indiana, USA

Correspondence

Steven W. Pipe, Department of Pediatrics and Communicable Diseases, Division of Hematology and Oncology, The University of Michigan Health System, 1500 East Medical Center Drive, D4207 MPB/5718, Ann Arbor, MI 48109, USA.

Email: ummdswp@med.umich.edu

Funding information

LFB SA

Abstract

Introduction: Eptacog beta is a new recombinant activated human factor VII bypassing agent approved in the United States for the treatment and control of bleeding in patients with haemophilia A or B with inhibitors 12 years of age or older.

Aim: To prospectively assess in a phase 3 clinical trial (PERSEPT 2) eptacog beta efficacy and safety for treatment of bleeding in children <12 years of age with haemophilia A or B with inhibitors.

Methods: Using a randomised crossover design, subjects received initial doses of 75 or 225 µg/kg eptacog beta followed by 75 µg/kg dosing at predefined intervals (as determined by clinical response) to treat bleeding episodes (BEs). Treatment success criteria included a haemostasis evaluation of 'excellent' or 'good' without use of additional eptacog beta, alternative haemostatic agent or blood product, and no increase in pain following the first 'excellent' or 'good' assessment.

Results: Treatment success proportions in 25 subjects (1–11 years) who experienced 546 mild or moderate BEs were 65% in the 75 µg/kg initial dose regimen (IDR) and 60% in the 225 µg/kg IDR 12 h following initial eptacog beta infusion. By 24 h, the treatment success proportions were 97% for the 75 µg/kg IDR and 98% for the 225 µg/kg IDR. No thrombotic events, allergic reactions, neutralising antibodies or treatment-related adverse events were reported.

Conclusion: Both 75 and 225 µg/kg eptacog beta IDRs provided safe and effective treatment and control of bleeding in children <12 years of age.

KEYWORDS

eptacog beta, haemophilia, inhibitors, PERSEPT, paediatric, recombinant FVIIa

1 | INTRODUCTION

Inhibitor development against factor VIII (FVIII) or factor IX (FIX) represents a serious complication in the management of haemophilia, resulting in compromised therapy effectiveness, increased morbidity and mortality, progressive joint disease and reduced quality of life.^{1–3} Inhibitors develop in about 20%–30% of patients with severe haemophilia A and up to 10% of patients with severe haemophilia B.^{4,5} Inhibitor eradication can be achieved through immune tolerance induction (ITI), but this approach is not effective in all patients (notably less success for haemophilia B patients with inhibitors).^{6,7}

During ITI or in the absence of inhibitor eradication, bypassing agents (BPAs) such as the recombinant activated human factor VII (rFVIIa) products eptacog beta (SEVENFACT®; HEMA Biologics, LLC and LFB SA)⁸ and eptacog alfa (NovoSeven® RT; Novo Nordisk),⁹ and the plasma-derived activated prothrombin complex concentrate (aPCC, FEIBA®; Takeda)¹⁰ are needed to manage bleeding episodes (BEs) in inhibitor patients. Emicizumab (Hemlibra®; Roche) is a humanised bispecific antibody with affinity for FIX/activated FIX and factor X (substituting for the cofactor activity of FVIII), and promotes effective haemostasis in patients with haemophilia A, even in the presence of inhibitors.¹¹ However, emicizumab is administered as a

prophylactic therapy and cannot treat breakthrough bleeding events. Thus, BPAs are still required for management of acute bleeds. rFVIIa (either eptacog beta or eptacog alfa) is the recommended treatment for breakthrough bleeds in haemophilia A patients with inhibitors who use emicizumab prophylaxis.¹²

Eptacog beta is a new rFVIIa BPA produced in a transgenic rabbit expression system.¹³ Approval of eptacog beta for treatment and control of BEs occurring in adult and adolescent haemophilia A or B patients with inhibitors (≥ 12 years of age) was obtained from the US Food and Drug Administration (FDA) in 2020.⁸ The pivotal phase 3 trial (PERSEPT 1; NCT02020369) demonstrated both 75 and 225 $\mu\text{g}/\text{kg}$ initial dose regimens (IDRs) of eptacog beta were effective in controlling mild or moderate BEs, with three severe BEs being successfully treated.¹⁴ At 12 h post-initial eptacog beta infusion, the reported treatment success proportions for mild or moderate BEs were 82% (75 $\mu\text{g}/\text{kg}$ IDR) and 91% (225 $\mu\text{g}/\text{kg}$ IDR).⁸ Following an initial 225 $\mu\text{g}/\text{kg}$ dose, 84% of mild or moderate BEs were controlled within 3 h and required no further dosing. A low rebleeding rate (1/465 mild or moderate BEs, 0.2%) was reported.¹⁴ The unpredictable inter- and intra-patient efficacy of existing BPAs¹⁵ has highlighted the need to further optimise treatment outcomes through continued development of new safe and efficacious BPAs such as eptacog beta.^{14,16,17}

The safety and efficacy of eptacog beta for BE treatment and control were further examined in a prospective phase 3 clinical trial (PERSEPT 2) enrolling paediatric subjects younger than 12 years of age with haemophilia A or B and inhibitors. PERSEPT 2 is the first reported prospective study to focus exclusively on this population for bleed treatment with a rFVIIa BPA. Results from this trial are described here.

2 | METHODS

2.1 | Eligibility criteria

Male subjects younger than 12 years of age with haemophilia A or B with inhibitors to FVIII or FIX respectively were eligible to enroll. Additional inclusion and exclusion criteria are listed in Table S1.

2.2 | Trial design

PERSEPT 2 was a multicentre, open-label, randomised phase 3 trial evaluating the safety, immunogenicity, pharmacokinetics (PK) and efficacy of two IDRs of eptacog beta for treatment of BEs (NCT02448680). PERSEPT 2 was designed as a crossover trial, with subject randomisation to either a 75 or a 225 $\mu\text{g}/\text{kg}$ IDR with crossover to the alternate IDR every 3 months without a washout period for the duration of the trial (replicating the trial design and dosing schedules of PERSEPT 1;¹⁴ Figure 1). Subjects received an initial infusion of 75 or 225 $\mu\text{g}/\text{kg}$ eptacog beta (per IDR randomisation) in a non-bleeding state for safety assessment and PK measurement purposes. Serum sample testing for anti-eptacog beta antibodies was performed as previously described.¹⁸

BEs were characterised as mild, moderate, or severe (Table S2). Subjects were advised to initiate treatment with a 2-min intravenous infusion of either 75 or 225 $\mu\text{g}/\text{kg}$ eptacog beta (depending upon IDR randomisation) as soon as possible after recognising bleeding symptoms. BE treatment was anticipated to occur in the home or community setting in most cases. Evaluations of efficacy and need for additional dosing were made at 3 and 9 h after initial infusion for subjects in the 75 and 225 $\mu\text{g}/\text{kg}$ IDRs, respectively. Need for additional 75 $\mu\text{g}/\text{kg}$ dosing was assessed every 3 h thereafter for subjects in both IDRs until the 21 h timepoint, with a final efficacy assessment at 24 h (Figure 1B). No additional study drug was permitted after 21 h in either IDR; if further treatment was required at 24 h, then alternative therapies could be initiated. Subjects who received at least three doses of eptacog beta while in the 75 $\mu\text{g}/\text{kg}$ IDR received the same cumulative amount of eptacog beta by 9 h post-initial infusion as subjects in the 225 $\mu\text{g}/\text{kg}$ IDR (Figure 1B). The 9 h interval between initial and subsequent doses of eptacog beta in the 225 $\mu\text{g}/\text{kg}$ IDR is supported by previous PK analyses.¹⁹ The protocol for treating severe BEs is described by Wang et al.¹⁴


Control of mild or moderate BEs following eptacog beta treatment was rated by the parent or caregiver according to a four-point haemostasis evaluation scale ('excellent', 'good', 'moderate' or 'none'; Table S3). Assessments took place in conjunction with the paediatric subject when possible, depending upon age and verbal abilities. Haemostasis evaluations for severe BEs were reported by the treating physician.

Treatment of a mild or moderate BE was considered successful if the following four criteria were met: (i) a haemostasis evaluation of 'excellent' or 'good' was obtained; (ii) no additional eptacog beta was given within 24 h after the first 'excellent' or 'good' response was noted; (iii) no alternative haemostatic agent or blood product was needed and (iv) pain associated with the BE did not increase following the initial 'excellent' or 'good' response. A Visual Analogue Scale (VAS) was used to rate pain on a scale of 0 (no pain) to 100 mm (worst possible pain), with subject or caregiver marking a position on a straight 100 mm line to represent relative pain severity.²⁰ The primary efficacy endpoint for PERSEPT 2 was defined as the successful treatment of mild or moderate BEs at 12 h following initial eptacog beta administration. The proportion of mild or moderate BEs successfully treated at 12 h was compared to an objective performance criterion (OPC) of 55%, which was derived from analysis of published studies examining rFVIIa efficacy in adult and paediatric subjects with haemophilia A or B and inhibitors, as previously described.¹⁴ As prospective clinical trials of bleed control with BPAs in children under 12 years of age are without precedent, such an OPC was considered the only benchmark available when the PERSEPT 2 trial was designed.

2.3 | Statistical analysis

PERSEPT 2 was designed to detect a true treatment success proportion of 0.70 for mild or moderate BEs with at least 80% power when comparing with the OPC of 0.55, assuming a one-sided asymptotically

(A)



(B)




FIGURE 1 (A) Subject dispositions for PERSEPT 2. (B) Treatment protocol for mild and moderate BEs in PERSEPT 2, with dosing schedules for 75 and 225 µg/kg IDRs indicated

normal test adjusted for multiplicity ($\alpha = 0.0125$) for each dosing regimen, correlated BEs, and 8 BEs per IDR per patient. Success proportion estimates at 12 and 24 h were based on observed cases reported and summarised using descriptive statistics. Estimates and test comparisons were based on an asymptotically normal estimator taking into account within-patient correlation. Statistical analyses were performed using SAS[®] 9.4.

2.4 | Ethics

The study protocol was reviewed and approved by institutional review boards or independent ethics committees at each study site, and was conducted in compliance with good clinical practice as described in the principles stated in the Declaration of Helsinki.²¹ Assent from the subjects and written informed consent from parents or legal guardians of the subjects were obtained at enrollment.

3 | RESULTS

3.1 | Subject population

Subject demographics are shown in Table 1. Twenty-five subjects were enrolled from 31 screened (Figure 1A). None were receiving BPA or emicizumab prophylaxis (PERSEPT 2 was completed prior to emicizumab regulatory approval). Four subjects discontinued PERSEPT 2 early, either by physician decision (two subjects: one was nonadherent and another was placed on prophylaxis) or due to withdrawal of consent (two subjects). No subject was discontinued due to an adverse event.¹⁸

3.2 | Efficacy

Subjects experienced 549 BEs: 546 BEs were mild or moderate and three were severe. Subjects experienced 239 mild or moderate BEs in

TABLE 1 PERSEPT 2 trial demographics

Subjects (N = 25)	Summary
Age, y	
Mean (SD)	4.9 (3.3)
Median	5.0
1st/3rd quartile	2/8
Minimum/maximum	1/11
Race, n (%)	
Black or African American	7 (28.0)
White	18 (72.0)
Ethnicity, n (%)	
Hispanic or Latino	3 (12.0)
Not Hispanic or Latino	22 (88.0)
Weight, kg	
Mean (SD)	20.9 (10.8)
Median	19.0
1st/3rd quartile	12.5/26.9
Minimum/maximum	8.2/52.0
Haemophilia type, n (%)	
Haemophilia A	23 (92.0)
Haemophilia B	2 (8.0)
Inhibitor titre, n (%) ^a	
BU ≥ 5	18 (72.0)
BU < 5 and high anamnestic response expected	6 (24.0)
BU < 5 and refractory to increased factor replacement dosing anticipated	1 (4.0)

SD, standard deviation; BU, Bethesda units.

^aInhibitor titer assessment performed at screening. Anamnestic and refractory response as indicated in disease history.

the 75 µg/kg IDR and 307 mild or moderate BEs in the 225 µg/kg IDR, with 92% of all mild or moderate BEs treated at home. The treatment success proportions and 95% confidence intervals (CIs) at 12 h for mild or moderate BEs were 65.4% (95% CI: [52.3%, 78.5%]) for the 75 µg/kg

IDR and 60.3% (95% CI: [48.2%, 72.3%]) for the 225 µg/kg IDR. At 24 h, the treatment success proportions were 97.4% (95% CI: [91.3%, 100.0%]) for the 75 µg/kg IDR and 98.0% (95% CI: [94.5%, 100.0%]) for the 225 µg/kg IDR (Figure 2). The difference in treatment success proportion between the two IDRs was not statistically significant at 12 or 24 h. Overall treatment success proportion for all mild or moderate BEs in both IDRs combined was 62.5% (95% CI: [51.1%, 74.0%]) at 12 h and 97.8% (95% CI: [93.1%, 100.0%]) at 24 h (Figure 2). Three subjects experienced a single severe BE and were treated using a severe BE dosing regimen (Table 2). Twelve (2.2%) of the mild or moderate BEs required an alternative haemostatic agent (either aPCC or eptacog alfa) for bleed control, with eight of these BEs occurring in a single subject.

Bleeding into joints accounted for 68% of the 546 mild or moderate BEs (Figure 3), and recurrent BEs in the same joint (a target joint) represented 19.9% of all joint BEs. The difference in treatment success proportions between target joint and non-target joint BEs was not statistically significant, for either IDR considered separately or for both IDRs combined. A median of two doses was required for bleed control in the 225 µg/kg IDR, and a median of three doses was required in the 75 µg/kg IDR. Eight (1.5%) of all mild or moderate BEs experienced a recurrence of bleeding (defined as bleeding in the same joint or anatomical location within 24 h after an initial 'good' or 'excellent' haemostasis evaluation). No assessments of BE recurrence were made beyond 24 h.

The overall proportion of BE treatments assessed as 'excellent' or 'good' at 12 h (64.3%; 95% CI: [52.6%, 76.1%]) and at 24 h (97.6%; 95% CI: [93.0%, 100%]) were similar to the success proportions calculated at 12 and 24 h when applying all four treatment success criteria. Successful pain relief was observed at 12 h after initial eptacog beta administration in 92.8% (75 µg/kg IDR) and 90.8% (225 µg/kg IDR) of mild or moderate BEs. Mean percent decreases in VAS pain score from baseline at 12 h after initial infusion were 70.9% (75 µg/kg IDR) and 64.5% (225 µg/kg IDR).

A 4-year-old subject with haemophilia A and a low body mass index (11.5 kg/m²) experienced 46 (8.4%) of the 546 mild or moderate BEs in PERSEPT 2, and had an outsized effect on efficacy results. This


**FIGURE 2** Success proportions and 95% confidence intervals for mild or moderate BE treatment at 12 and 24 h

TABLE 2 Severe BEs in PERSEPT 2

Subject	Age (y)	Haemophilia type	Severe BE type	Severe BE dosing regimen ($\mu\text{g}/\text{kg}$)	Hospital visit	Haemostasis evaluation (12 h)	Haemostasis evaluation (24 h)
Subject 1	9	Haemophilia A	Spontaneous renal haemorrhage ^a	225	Yes	Moderate	Moderate
Subject 2	8	Haemophilia A	Traumatic intracranial bleed ^b	225 ^c	Yes	None	None
Subject 3	6	Haemophilia B	Traumatic left elbow bleed	225 ^c	No	Not recorded	Not recorded

BE, bleeding episode.

^aSubject- and physician-reported haemostasis evaluations were both 'good' after approximately 4 days of treatment with eptacog beta, a treatment duration consistent with consensus management guidelines.²⁹

^bThe intracranial bleed (confirmed by computed tomography [CT] scan) was caused by head trauma and resolved with treatment. Subject 2 received eptacog beta for 3 days followed by aPCC (FEIBA[®]) for 17 days, a treatment duration that is consistent with previously-described intracranial haemorrhage management for haemophilia A or B patients with inhibitors³⁰ as well as with consensus treatment guidelines.²⁹ CT scans on days 2, 4 and 5 following BE onset showed no further intracranial bleeding.

^cThe 225 $\mu\text{g}/\text{kg}$ severe BE dosing regimen (as detailed by Wang et al.¹⁴) was not followed: either scheduled eptacog beta infusions were delayed by as long as 3 h (for the left elbow BE) or eptacog beta amounts and dosing intervals other than those specified by the protocol were used (for the intracranial bleed).




FIGURE 3 Anatomical distributions of the 546 mild or moderate BEs in PERSEPT 2. The percentage for each anatomical site of the total number of mild or moderate BEs is indicated

subject exhibited a delayed response to eptacog beta treatment, with a treatment success proportion at 12 h of 5.6% for the 75 $\mu\text{g}/\text{kg}$ IDR and 14.3% for the 225 $\mu\text{g}/\text{kg}$ IDR. Treatment success proportions at 24 h for both IDRs were 100%. Serum samples from this subject tested negative for anti-eptacog beta antibodies. When excluding this outlier subject from the PERSEPT 2 analyses, the treatment success proportions and 95% CIs at 12 h for mild or moderate BEs increase to 70.3% (95% CI: [59.8%, 80.8%]) for the 75 $\mu\text{g}/\text{kg}$ IDR and 65.0% (95% CI: [54.5%, 75.4%]) for the 225 $\mu\text{g}/\text{kg}$ IDR.

3.3 | Safety

As previously described, eptacog beta was well tolerated: no thrombotic events, allergic, hypersensitivity or anaphylactic reactions, neutralising anti-eptacog beta antibodies or treatment-related adverse events were observed.¹⁸ Three serious adverse events (paresis, intracranial bleed and dysentery, all resolving with treatment) were assessed as unrelated to eptacog beta administration.¹⁸

4 | DISCUSSION

In the current study, the safety and efficacy of a new rFVIIa BPA (eptacog beta) for BE treatment was examined: 549 BEs in 25 paediatric subjects with haemophilia A or B and inhibitors under 12 years of age were evaluated. As with the study in adults and adolescents (PERSEPT 1),¹⁴ a four-part composite of haemostasis and pain criteria (see Section 2) was used to determine treatment success in PERSEPT 2. This extensive set of treatment success criteria provides confidence that satisfactory clinical benefits were received by study subjects, acknowledging the subjective nature of haemostasis evaluations and pain assessment in young children.

The subjective aspects of determining bleeding control are well-recognised challenges in assessing treatment efficacy in haemophilia patients, particularly in paediatric subjects.²² Decisions to cease or continue treating BEs in children with haemophilia have a significant subjective component^{23,24} and are frequently made indirectly by the caregiver, as younger patients may not be fully capable of

identifying and communicating when bleeds have resolved. Caregivers report uncertainty in ascertaining exactly when BEs have resolved, and may base continued treatment on rebleeding concerns or on prior experience with similar bleeds.²³ Such factors may contribute to the longer rFVIIa treatment duration seen in children over that observed in adults.²⁵ The subjective aspects of evaluating pain in children²⁰ further complicate haemostasis assessments made by caregivers.

The treatment success proportion for eptacog beta at 12 h was compared to an OPC (55%) derived from published rFVIIa clinical studies including adult and paediatric subjects with haemophilia A or B and inhibitors. This OPC had been previously used in the PERSEPT 1 trial as a benchmark for eptacog beta treatment success in treatment-experienced adults and adolescents,¹⁴ and was the only primary endpoint comparator available at the time the PERSEPT 2 trial was designed. An appropriate primary efficacy threshold specifically for prospective paediatric haemophilia trials such as PERSEPT 2 has never been determined.

The overall treatment success proportion of 546 mild or moderate BEs in PERSEPT 2 was 63% at 12 h (Figure 2). As the lower bound of the 95% CI for this point estimate did not exceed 55%, the treatment success proportion at 12 h was not significantly different from the OPC. Similarly, the bleed treatment success proportions at 12 h of 65% for the 75 µg/kg IDR and 60% for the 225 µg/kg IDR (Figure 2) did not significantly differ from the OPC. While the 12-h primary endpoint and the OPC were not statistically different, by 24 h the bleed treatment success proportions were much higher: the overall treatment success was 98% at 24 h (Figure 2), with bleed treatment success proportions of 97% for the 75 µg/kg IDR and 98% for the 225 µg/kg IDR (Figure 2). Several other trial findings support the efficacy of eptacog beta in PERSEPT 2 subjects: (i) only 1.5% of all mild or moderate BEs recurred at the same anatomical site within 24 h of the first 'excellent' or 'good' haemostasis evaluation; (ii) bleed control was achieved with a median of two doses in the 225 µg/kg IDR and three doses in the 75 µg/kg IDR for mild or moderate BEs; (iii) only 2.2% of all BEs required alternative treatment and (iv) by 12 h after initial eptacog beta infusion, mean VAS pain scores were reduced by 70.9% (75 µg/kg IDR) and 64.5% (225 µg/kg IDR) from baseline values at BE onset. No thrombotic events, allergic, hypersensitivity, or anaphylactic reactions, neutralising anti-eptacog beta antibodies or treatment-related adverse events were reported. The totality of these data indicate that eptacog beta was safe and effective by 24 h in treating and controlling mild or moderate bleeding in the study participants. As BE resolution was achieved using fewer median number of infusions in the 225 µg/kg IDR as compared to the 75 µg/kg IDR, the 225 µg/kg IDR may potentially be more attractive for patients and less burdensome to caregivers.

Previous pharmacokinetic and laboratory pharmacodynamics analyses have demonstrated peak eptacog beta plasma levels (C_{max}), peak thrombin generation, and clot firmness exhibit a dose-dependent relationship with eptacog beta.¹⁹ In accord with these findings, clinical trial data from adults and adolescents in PERSEPT 1 showed a higher treatment success proportion in the 225 µg/kg IDR as compared to the 75 µg/kg IDR at 12 h (91% and 82% in the 225 and 75 µg/kg IDRs, respectively),⁸ a result consistent with a dose-dependent throm-

bin burst driving haemostasis at the site of injury. A similar outcome in PERSEPT 2 might be reasonably anticipated; however, observed efficacies for the two IDRs at 12 h were instead comparable in magnitude (60% and 65% in the 225 and 75 µg/kg IDRs, respectively), and lower overall than those seen in PERSEPT 1. An elevated weight-adjusted clearance of eptacog beta in paediatric subjects (as previously reported for eptacog alfa²⁶⁻²⁸) provides a plausible explanation for the observed 12-h haemostatic response (Figure 2); however, the comparable efficacies observed for the two IDRs at 12 h might not be consistent with such a model being the sole factor for the observed results. Assuming similar weight-adjusted eptacog beta clearance for subjects in either IDR, any increased clearance in paediatric subjects should impact treatment success proportions for both IDRs to the same degree, preserving the same higher treatment success proportion in the 225 µg/kg IDR relative to the 75 µg/kg IDR seen in PERSEPT 1. The observed haemostatic response might be better explained by the subjective nature that surrounds the determination of BE resolution by caregivers, as well as a bias towards continued treatment out of rebleeding concerns^{23,24} (regardless of IDR). A lack of caregiver clarity regarding BE cessation along with a bias towards retreatment would contribute towards conservative estimates of treatment success proportions at the 12-h timepoint, and could effectively mask any real differences between treatment success proportions for the two IDRs at 12 h.

Additional factors further inform interpretation of trial results. While a limited number of subjects participated in PERSEPT 2 (25 subjects), haemostatic efficacy was reported as the proportion of successfully-treated BEs and a substantial number of mild or moderate BEs were available for analysis (546 BEs). A 4-year-old subject with haemophilia A who experienced 46 of the 546 mild or moderate BEs and a delayed response to eptacog beta exerted an outsized effect on efficacy results, resulting in a reduced treatment success proportion for each IDR at 12 h after initial BE treatment. The treatment success proportions for these 46 BEs at 24 h were 100% for both IDRs, demonstrating the effectiveness of eptacog beta for control of bleeding in this subject by the 24-h timepoint.

5 | CONCLUSION

PERSEPT 2 is the first prospective study of bleed treatment with a rFVIIa BPA focused solely on interval analysis of clinical response in haemophilia A or B subjects with inhibitors younger than 12 years of age. Both eptacog beta IDRs (75 and 225 µg/kg) provided safe and effective treatment and control of bleeding by 24 h to the trial subjects: no thrombotic events, allergic reactions or treatment-related adverse events were reported; a significant proportion of BEs were successfully treated at 12 h; and nearly all BEs were resolved at 24 h. As such, eptacog beta potentially offers an important therapeutic option to patients, caregivers and health care providers for BE treatment.

ACKNOWLEDGEMENTS

The Program for the Evaluation of Recombinant Factor Seven Efficacy by Prospective Clinical Trial - PERSEPT was funded by LFB SA. The

authors would like to thank the participants of the PERSEPT 2 trial and their families, without whom none of this work would be possible. The authors also wish to thank the investigators at each participating study site, as well as Sonia Nasr, PhD for assistance during the editing of this manuscript.

DISCLOSURES

S.W.P. has received consulting fees from Apcintex, ASC Therapeutics, Bayer, BioMarin, Catalyst Biosciences, CSL Behring, GenVentiv, HEMA Biologics, Freeline, LFB, Novo Nordisk, Pfizer, Regeneron/Intellia, Roche/Genentech, Sangamo Therapeutics, Sanofi, Takeda, Spark Therapeutics and uniQure. M. Chitlur has received research funding from Genentech and Agios Pharmaceuticals, and has acted as a paid consultant and received honoraria for participation in advisory boards for Novo Nordisk, Takeda, CSL Behring, Genentech, Bio Products Laboratories and Agios Pharmaceuticals. M. Carcao has received research support from Bayer, Bioverativ/Sanofi, CSL Behring, Novo Nordisk, Octapharma, Pfizer, Roche, and Shire/Takeda; has received honoraria for speaking/participating in advisory boards from Bayer, Bioverativ/Sanofi, CSL Behring, Grifols, LFB, Novo Nordisk, Pfizer, Roche, and Shire/Takeda; and has no stocks or equity in any pharmaceutical company, and is not part of any company board. G.C. has received honoraria as a speaker/participant on advisory boards for Alexion, Bayer, Shire/Takeda, CSL Behring, Novo Nordisk, Sobi, Roche, uniQure, Sanofi, Werfen, Kedrion, LFB and Grifols. J.A.D. has acted as a speaker for Genentech, Sanofi Genzyme, and BioMarin, and as a consultant for Spark, Takeda and Novo Nordisk. J.D. has acted as a consultant for Bayer and HEMA Biologics, and has been on the speaker's bureau for Bayer. A.L.D. has acted as a paid consultant for Genentech, Kedrion, CSL Behring and BioMarin; has received research funding from Takeda, BioMarin, Freeline, Novo Nordisk, Sanofi and the American Society of Thrombosis and Hemostasis; and is on the board of World Federation of Hemophilia USA. M.E. has received honoraria for consulting and participating in advisory boards from BioMarin, Novo Nordisk, Genentech, Sanofi, Takeda, Pfizer, Kedrion, CSL Behring and NHF. O.K. has served as a consultant or medical advisor and has been reimbursed for services by Genentech, Octapharma, BioMarin, CSL Behring, HEMA Biologics, Kedrion, Takeda and Novo Nordisk. J.M. has received research grants from Bayer, Biogen, BioMarin, CSL, Novo Nordisk, Sobi, Roche and uniQure; has served on scientific advisory committees of Amgen, Bayer, Biotest, Biogen, Baxalta, CSL Behring, Catalyst Biosciences, Novo Nordisk, Roche and Spark; and has been a member of the speaker bureau of Alnylam, Bayer, Biotest, Biogen, Novo Nordisk, Pfizer, Sobi, Shire, Roche, ISTH and WFH. S.L.M. has been on advisory boards for Novo Nordisk, Shire, Genentech, Sanofi and BioMarin; is a paid consultant for Spark and Pfizer; and receives investigator-initiated research funding as a Co-I from Octapharma and Genentech. C.N. has received honoraria/consultation fees or grants/research support from Bayer, CSL Behring, Freeline, LFB, Novo Nordisk, Octapharma, Pfizer, Roche-Chugai, Sanofi, Shire-Takeda, Sobi and Spark Therapeutics. M.R.'s employers have received research funding from Bayer, BioMarin, CSL Behring, Genentech, Grifols, HEMA Biologics, LFB, Novo Nordisk, Octapharma, Pfizer, Sanofi, Spark, Takeda and uniQure.

M.R. has acted as a paid consultant to Catalyst Biosciences, CSL Behring, Genentech, HEMA Biologics, Kedrion, Novo Nordisk, Pfizer, Sanofi, Takeda and uniQure. M.R. is on the board of directors of Foundation for Women and Girls with Blood Disorders and Partners in Bleeding Disorders. M.R. is an employee of the American Thrombosis and Hemostasis Network and Oregon Health & Science University. T.C-S. has served as a consultant and scientific advisory board member for Octapharma, Bayer, Novo Nordisk, CSL Behring, Genentech, BioMarin, Takeda, HEMA Biologics, Kedrion and Pfizer; and has served on a speakers bureau for Octapharma, Genentech, BioMarin, Novo Nordisk, CSL Behring, Takeda, Grifols, and Spark. T.C-S. has received honoraria from Octapharma, Genentech, CSL Behring, Novo Nordisk, Takeda, BPL, BioMarin, Grifols and Pfizer; and has received research funding from Spark, BioMarin and Pfizer. M.W. has been a consultant and/or advisor to Bioverativ/Sanofi, Takeda, CSL Behring, Catalyst Biosciences, Novo Nordisk, Bayer, Octapharma, Genentech, HEMA Biologics, BioMarin and uniQure, and was a study investigator for HEMA Biologics for research carried out in this work. J.W. has received grant support from Alnylam, Baxalta, Novo Nordisk, Octapharma, Rigel Pharmaceuticals, Roche, Shire/Takeda and Sobi; and has received honoraria from Alexion, Baxalta, CSL Behring, Ferring Pharmaceuticals, LFB, Novo Nordisk, Octapharma, Roche, Sanofi/Genzyme, Shire/Takeda, Siemens, Sobi and Werfen. G.Y. has received honoraria for consulting for Genentech/Roche and a grant from Genentech. G.Y. also has received honoraria from BioMarin, Grifols, Pfizer, Sanofi, Spark, and Takeda; and has grants from Grifols and Takeda. W.A.A. works as a consultant for HEMA Biologics, LLC, and has received fees for speaking and consulting. D.B. is an employee of LFB-USA. C.M. and I.S.M. are employees of HEMA Biologics, LLC, and I.S.M. was formerly a consultant with HEMA Biologics, LLC. E.S. is an employee of LFB. T.A.W. is an analyst/medical writer for GLOVAL, LLC. A.D.S. has served on advisory boards or as consultant for Genentech, Roche, Novo Nordisk, BioMarin, Bioverativ, Sanofi, ProMetric Bio Sciences, Sangamo, Sigilon, and Takeda; and has received research funding from these organisations plus Agios, OPKO Global Bio Therapeutics, Kedrion, Octapharma, and Novartis. C.H., J.J., I.H.M., U.N-G., O.S., K.V.V. and L.V.M. have no competing interests to declare.

AUTHOR CONTRIBUTIONS

All of the authors analyzed and interpreted the data. Thomas A. Wilkinson developed the initial draft, and all authors provided critical input and edited the manuscript. All authors reviewed and approved the final submission.

DATA AVAILABILITY STATEMENT

Data available from the authors upon reasonable request.

ORCID

Steven W. Pipe  <https://orcid.org/0000-0003-2558-2089>

Cédric Hermans  <https://orcid.org/0000-0001-5429-8437>

Manuel Carcao  <https://orcid.org/0000-0001-5350-1763>

Giancarlo Castaman  <https://orcid.org/0000-0003-4973-1317>

Amy L. Dunn  <https://orcid.org/0000-0002-9645-6365>

Miguel Escobar  <https://orcid.org/0000-0002-2944-0240>
 Johnny Mahlangu  <https://orcid.org/0000-0001-5781-7669>
 Shannon L. Meeks  <https://orcid.org/0000-0002-3683-8644>
 Laura Villarreal Martinez  <https://orcid.org/0000-0003-0313-7312>
 Michael Wang  <https://orcid.org/0000-0001-9289-4862>
 Guy Young  <https://orcid.org/0000-0001-6013-1254>

REFERENCES

- Scalone L, Mantovani LG, Mannucci PM, Gringeri A, COCIS Study Investigators. Quality of life is associated to the orthopaedic status in haemophilic patients with inhibitors. *Haemophilia*. 2006;12(2):154-162.
- Neufeld EJ, Recht M, Sabio H, et al. Effect of acute bleeding on daily quality of life assessments in patients with congenital hemophilia with inhibitors and their families: observations from the dosing observational study in hemophilia. *Value Health*. 2012;15(6):916-925.
- Walsh CE, Jimenez-Yuste V, Auerswald G, Grancha S. The burden of inhibitors in haemophilia patients. *Thromb Haemost*. 2016;116 (Suppl. 1):S10-17.
- Astermark J. Overview of inhibitors. *Semin Hematol*. 2006;43(Suppl 4):S3-7.
- Male C, Andersson NG, Rafowicz A, et al. Inhibitor incidence in an unselected cohort of previously untreated patients with severe haemophilia B: a PedNet study. *Haematologica*. 2021;106(1):123-129.
- Ljung R, Auerswald G, Benson G, et al. Inhibitors in haemophilia A and B: management of bleeds, inhibitor eradication and strategies for difficult-to-treat patients. *Eur J Haematol*. 2019;102(2):111-122.
- Astermark J, Holstein K, Abajas YL, et al. The B-natural study-The outcome of immune tolerance induction therapy in patients with severe haemophilia B. *Haemophilia*. 2021;27(5):802-813.
- SEVENFACT® [Package Insert]. Louisville, KY: HEMA Biologics, LLC; 2020.
- Hedner U. Recombinant activated factor VII: 30 years of research and innovation. *Blood Rev*. 2015;29(Suppl 1):S4-8.
- Négrier C, Gomperts ED, Oldenburg J. The history of FEIBA: a lifetime of success in the treatment of haemophilia complicated by an inhibitor. *Haemophilia*. 2006;12(s5):4-13.
- Oldenburg J, Mahlangu JN, Kim B, et al. Efficacy of Emicizumab Prophylaxis in Hemophilia A with Inhibitors. *N Engl J Med*. 2017;377(9):809-818.
- Medical and Scientific Advisory Council (MASAC). *MASAC Recommendations Concerning Products Licensed for the Treatment of Hemophilia and Other Bleeding Disorders*. MASAC Document #263. National Hemophilia Foundation; 2020.
- Chevreux G, Tilly N, Leblanc Y, et al. Biochemical characterization of LR769, a new recombinant factor VIIa bypassing agent produced in the milk of transgenic rabbits. *Haemophilia*. 2017;23(4):e324-e334.
- Wang M, Lawrence JB, Quon DV, et al. PERSEPT 1: a phase 3 trial of activated eptacog beta for on-demand treatment of haemophilia inhibitor-related bleeding. *Haemophilia*. 2017;23(6):832-843.
- Astermark J, Donfield SM, DiMichele DM, et al. A randomized comparison of bypassing agents in hemophilia complicated by an inhibitor: the FEIBA NovoSeven Comparative (FENOC) Study. *Blood*. 2007;109(2):546-551.
- Meeks SL, Leissing CA. The evolution of factor VIIa in the treatment of bleeding in haemophilia with inhibitors. *Haemophilia*. 2019;25(6):911-918.
- Shapiro AD, Mitchell IS, Nasr S. The future of bypassing agents for hemophilia with inhibitors in the era of novel agents. *J Thromb Haemost*. 2018;16(12):2362-2374.
- Escobar M, Castaman G, Boix SB, et al. The safety of activated eptacog beta in the management of bleeding episodes and perioperative haemostasis in adult and paediatric haemophilia patients with inhibitors. *Haemophilia*. 2021;27(6):921-931.
- Ducore J, Lawrence JB, Simpson M, et al. Safety and dose-dependency of eptacog beta (activated) in a dose escalation study of non-bleeding congenital haemophilia A or B patients, with or without inhibitors. *Haemophilia*. 2017;23(6):844-851.
- Humphries TJ, Kessler CM. The challenge of pain evaluation in haemophilia: can pain evaluation and quantification be improved by using pain instruments from other clinical situations? *Haemophilia*. 2013;19(2):181-187.
- World Medical Association. World Medical Association Declaration of Helsinki: ethical principles for medical research involving human subjects. *JAMA*. 2013;310(20):2191-2194.
- Amby LK, Seremetis S, Obergfell A, Bjerre J. Challenges of defining reliable clinical surrogate end points in haemophilia trials: a critical review. *Blood Coagul Fibrinolysis*. 2009;20(7):488-493.
- Valentino LA, Walsh CE, Reding MT, Young GA, Levendoglu-Tugal O, Cooper DL. Patient- and caregiver-reported bleeding symptoms and reasons for starting and stopping treatment with recombinant factor VIIa: analysis of the Dosing Observational Study in Haemophilia (DOSE). *Haemophilia*. 2012;18(4):554-560.
- Young G, Shapiro AD, Walsh CE, Gruppo RA, Gut RZ, Cooper DL. Patient/caregiver-reported recombinant factor VIIa (rFVIIa) dosing: home treatment of acute bleeds in the Dosing Observational Study in Hemophilia (DOSE). *Haemophilia*. 2012;18(3):392-399.
- Gruppo RA, Kessler CM, Neufeld EJ, Cooper DL. Assessment of individual dose utilization vs. physician prescribing recommendations for recombinant activated factor VII (rFVIIa) in paediatric and adult patients with congenital haemophilia and alloantibody inhibitors (CHwl): the Dosing Observational Study in Hemophilia (DOSE). *Haemophilia*. 2013;19(4):524-532.
- Santagostino E, Morfini M, Rocino A, Baudo F, Scaraggi F, Gringeri A. Relationship between factor VII activity and clinical efficacy of recombinant factor VIIa given by continuous infusion to patients with factor VIII inhibitors. *Thromb Haemost*. 2001;86(10):954-958.
- Villar A, Aronis S, Morfini M, et al. Pharmacokinetics of activated recombinant coagulation factor VII (NovoSeven) in children vs. adults with haemophilia A. *Haemophilia*. 2004;10(4):352-359.
- Klitgaard T, Nielsen TG. Overview of the human pharmacokinetics of recombinant activated factor VII. *Br J Clin Pharmacol*. 2008;65(1):3-11.
- Srivastava A, Santagostino E, Dougall A, et al. WFH Guidelines for the Management of Hemophilia, 3rd edition. *Haemophilia*. 2020;26(Suppl 6):1-158.
- Zanon E, Pasca S. Intracranial haemorrhage in children and adults with haemophilia A and B: a literature review of the last 20 years. *Blood Transfus*. 2019;17(5):378-384.

SUPPORTING INFORMATION

Additional supporting information may be found in the online version of the article at the publisher's website.

How to cite this article: Pipe SW, Hermans C, Chitlur M, et al. Eptacog beta efficacy and safety in the treatment and control of bleeding in paediatric subjects (<12 years) with haemophilia A or B with inhibitors. *Haemophilia*. 2022;1-9. <https://doi.org/10.1111/hae.14563>